

# **CHEMISTRY OF THIONITROSYL GROUP AND ITS COMPLEXES WITH RHODIUM (III)**

**A Thesis Submitted  
In Partial Fulfilment of the Requirements  
for the Degree of  
MASTER OF PHILOSOPHY**

**By  
D. KRISHNA MURTY RAJU**

**to the  
DEPARTMENT OF CHEMISTRY  
INDIAN INSTITUTE OF TECHNOLOGY, KANPUR  
APRIL, 1980**

RECEIVED  
LIBRARY  
20 NOV 1980

✓CHM-1880-M-RAJ-CHE

U. T. KANPUR  
CENTRAL LIBRARY

Acc. No. **A 63780**

20 NOV 1980

TO MY  
PARENTS & TEACHERS

### STATEMENT

I hereby declare that the matter embodied in this thesis is the result of investigations carried out by me in the Department of Chemistry, Indian Institute of Technology, Kanpur, India under the supervision of Professor U.C. Agarwala.

In keeping with the general practice of reporting scientific observations due acknowledgement has been made wherever the work described is based on the findings of other investigators.

*D. Krishnamurty Raju*

D. Krishnamurty Raju



Department of Chemistry  
Indian Institute of Technology, Kanpur

CERTIFICATE I

This is to certify that Mr. D. Krishna Murty Raju has satisfactorily completed all the courses required for the M.Phil. degree programme. These courses include:


Chem 501	Advanced Organic Chemistry I
Chem 521	Chemical Binding
Chem 524	Modern Physical Methods in Chemistry
Chem 541	Advanced Inorganic Chemistry I
Chem 542	Advanced Inorganic Chemistry II
Chem 581	Basic Biological Chemistry & Molecular Biology
Chem 800	General Seminar
Chem 801	Graduate Seminars
Chem 900	Post-Graduate Research



(S. Ranganathan)  
Professor and Head,  
Department of Chemistry



(D.N. Dhar)  
Convener  
Departmental Post-Graduate Committee



CERTIFICATE II

Certified that the work contained in this thesis,  
entitled "Chemistry of Thionitrosyl Group and its Complexes  
with Rhodium(III)" has been carried out by D.Krishna Murty  
Raju under my supervision and the same has not been submitted  
elsewhere for a degree.

*U.C. Agarwala*  
U.C. Agarwala  
Thesis Supervisor

Kanpur

April, 1980

## ACKNOWLEDGEMENT

It gives me great pleasure to place on recording deepsense of gratitude to Professor U.C. Agarwala for suggesting the problem and his inspiring guidance through out the course of the present investigations. I am grateful for his keenness, cooperation and affection towards me through out the project work.

I wish to express my sincere thanks to Mr. K.K. Pandey for his assistance and the fruitful and thought provoking discussions which I had with him many times.

Thanks are also due to my colleagues Dr. T. Singh, Mr. R. Parashad, Mr. S. Datta, Mr. S.K.S. Yadav, Mr. K.C. Jain, Mr. R. Shaheb, Mr. Arulsamy and other friends for their cooperation.

I am very much grateful to the management of D.N.R. College, Bhimavaram for their encouragement for research and sponsorship for the F.I.P.

Last but not least, I wish to thank Mr. Anil Kumar for his excellent typing of the manuscripts.

Kanpur  
April, 1980

D. Krishna Murty Raju

## CONTENTS

	page
STATEMENT	i
CERTIFICATE I	ii
CERTIFICATE II	iii
ACKNOWLEDGEMENT	iv
CHAPTER I	
CHEMISTRY OF THIONITROSYL GROUP	
INTRODUCTION	1
IONIC COMPOUNDS	20
COVALENT COMPOUNDS	26
METALLIC AND NONMETALLIC ADDUCTS AND COMPOUNDS OF SULFURNITRIDE	49
TRANSITION METAL THIONITROSYL COMPLEXES	53
REFERENCES	74
CHAPTER II	
THIONITROSYL COMPLEXES OF RHODIUM(III)	92
REFERENCES	100

.....

## CHAPTER I

### CHEMISTRY OF THIONITROSYL GROUP

#### INTRODUCTION

Thionitrosyl radical with one unpaired electron unlike its homolog, viz. NO, polymerizes so readily that it is not possible to isolate it as monomer solid or liquid and even in the gaseous phase, it has only a transient existence [1]. As a result of this, the studies pertaining to the structure, bonding and the reactivity of thionitrosyl\* have always been challenging and despite the large interest in the chemistry of thionitrosyl, the amount of work in this field is relatively much less and fragmentary.

Although several important reviews of the compounds containing sulfur and nitrogen [1-13], covering the chemistry of polymeric sulfur nitrides and those of sulfur-nitrogen-fluorine [14-20] have appeared in the literature, but there has been no comprehensive review exclusively on thionitrosyls. In the present review, our main aim has been to provide a comprehensive account to December 1979 of syntheses, physical properties, chemical reactions and structures of thionitrosyl, its ions and the transition metal thionitrosyl complexes.

---

\* In this review the term 'Thionitrosyl' is employed as a general name for the compounds of the NS group.

(a) Syntheses and General Properties

The thionitrosyl radical, like NO, H, I NO<sub>2</sub> and ClO<sub>2</sub> can also exist as NS<sup>+</sup> or NS<sup>-</sup> formed respectively by loosing or gaining an electron and therefore most of its properties should be approximately similar to those of NO. The presence of NS was observed for the first time by Fowler and Barrow in 1932 while studying the band spectrum of the emitted light after passing an electric discharge through a mixture of nitrogen and sulfur vapors. They observed a resemblance of the band spectrum with that of NO [21] and concluded the presence of NS in the mixture. A few other methods for its formation have since then been developed. These are:

- (i) By passing an electric discharge through SF<sub>6</sub> or sulfur vapour and nitrogen [21,22,23,24]
- (ii) By the interaction of sulfur vapour, H<sub>2</sub>S or SCl<sub>2</sub> and active nitrogen [25,29,26,22,27]
- (iii) By photolysing a mixture of COS and NF<sub>3</sub> 28
- (iv) By volatilization of (SN)<sub>x</sub> [30]
- (v) By thermal pyrolysis of S<sub>4</sub>N<sub>4</sub> over quartz wool and silver wool at a temperature greater than 300°C [31]

Evidence for the formation of SN<sup>+</sup> and SN<sup>-</sup> ions was provided for the first time by Dressel [32] and O'Hare [33] respectively during their spectroscopic studies. Subsequent work by other workers [34-37] led to the isolation of a

number of ionic salts containing  $\text{NS}^+$  ion which will be discussed at a later stage in the review.

NS bond length (149.7 pm) calculated from its spectroscopic moment of inertia [38] is shorter by 24-28 pm than the sum of the single bond covalent radii of nitrogen and oxygen. It suggests the bond order of NS between two and three which is parallel to that of NO because of the similar electronic structures of NS and NO molecules (Three bonding MO fully occupied and an odd electron in the antibonding MO).

The ionization potential for the process,  $\text{NS}(^2\Pi) \longrightarrow \text{NS}^+(^1\Sigma) + e^-$  at  $0^\circ\text{K}$  is 9.85 e.v. [39]. This value is slightly greater or similar to that for  $\text{NO} \longrightarrow \text{NO}^+ + e^-$  (9.8 e.v.) [40-41] and is lesser than that for  $\text{N}_2 \longrightarrow \text{N}_2^+ + e^-$  (15.5-17.2 e.v.)\* [42] or that for  $\text{O}_2 \longrightarrow \text{O}_2^+ + e^-$  (12.2-18.2 (12.2-18.2 e.v.)\* [43]. The lower value of the ionization potential of NS than that for  $\text{N}_2$  or  $\text{O}_2$  and similar to that of NO reflects the loss of an odd electron from the antibonding MO of NS or NO to produce more stable ion which is isoelectronic

\* The range in the value of the ionization potentials of  $\text{N}_2$  and  $\text{O}_2$  is given. The exact value will depend on the electronic state of the ion ( $\text{N}_2^+$  or  $\text{O}_2^+$ ).

Table I Thermodynamic Data of  $\text{NS}^+$ , NS and  $\text{NS}^-$  [61-69]

	Total energy hartree	Ionization Potential ev	Dissociation Energy(D) ev	Dipole Moment( $\mu$ )	Quadrupole Moment $Q \times 10^{-26}$ esu $\text{cm}^2$	$\Delta H_f$ ev
$\text{NS}^+$	-451.574631	24.32	$5.31 \pm 0.31$	3.893	0.1611	12.76
NS	-451.932861	9.85	$5.2 \pm 0.2$	1.732	-0.6734	2.91
$\text{NS}^-$	-451.95446	-	$4.05 \pm 0.2$	0.09	-3.89	-



to CO or N<sub>2</sub>. The ionic character of the NS bond (24%) calculated on the basis of Pauling's criterion [44] may be compared with that of PS bond (42%). This decrease in the ionicity of the bond parallels the difference between the electronegativities of phosphorus and nitrogen.

### (b) Thermodynamic Properties

The thermodynamic properties of NS, NS<sup>+</sup> and NS<sup>-</sup> have been studied both theoretically and from the spectral and thermochemical data by O'Hare [39]. The data relating to these properties are given in Table 1.

The dissociation energy of NS molecule was determined using Hildenbrand and Murad relationship [45] (4.8 e.v.) which is in excellent agreement with the one obtained by the method proposed by Gaydon [46] (4.79 e.v.). Equation,  $D_o^o(NS^+) = D_o^o(NS) + I(S) - I(NS)$  was subsequently applied to calculate the heat of dissociation of NS<sup>+</sup> (5.31 e.v.).

$D_o^o/D_{LBX}^o = \frac{5.256}{r_e I(S) - E_a(N)} + 0.448$ , where  $D_o^o$  is the true dissociation energy;  $D_{LBX}^o$ , dissociation energy determined by the Linear Birge-Sponer extrapolation of spectroscopic data method 53 ;  $r_e$  equilibrium internuclear distance (1.4957 Å for NS);  $I(S)$ , ionization energy of sulfur (10.36 e.v.) and  $E_a(N)$ , electron affinity of nitrogen (0.04 e.v.) 54

Dissociation of  $\text{NS}^-$  ( $^3\Sigma$ ) to give  $\text{N}$  ( $^4\text{S}$ ) and  $\text{S}^-$  ( $^2\text{P}$ ) involves the uncoupling of the same number of electron pairs as those in the dissociation of  $\text{NO}^-$  ( $^3\Sigma$ ) to  $\text{N}$  ( $^4\text{S}$ ) and  $\text{O}^-$  ( $^2\text{P}$ ). Assuming the molecular extra correlation energy,  $\Delta E_{\text{cor}}$ , to be the same for both the processes, the dissociation energy,  $D(\text{NA})$  and the electron affinity of  $\text{NA}$  and  $\text{A}$  (where  $\text{A}$  is  $\text{O}$ , or  $\text{S}$ ) were related as  $D(\text{NA}^-) = D(\text{NA}) + E_{\text{a}}(\text{NA}) - E_{\text{a}}(\text{A})$ . Using the data of  $\text{NO}$  [47-52],  $\Delta E_{\text{cor}}$  was calculated for  $\text{NO}^-$  which was also assumed to be the same for  $\text{NS}^-$  (0.1333 e.v.). Combination of the Hartree-Fock energies of  $\text{N}$  ( $^4\text{S}$ ),  $\text{S}^-$  ( $^2\text{P}$ ) and  $\text{NS}^-$ , yielded  $D(\text{NS})$  from which the value of  $D(\text{NS}^-)$  ( $4.05 \pm 0.2$  e.v.) was calculated after the addition of  $\Delta E_{\text{cor}}(\text{NS})$  to  $D(\text{NS})$ .

Using similar relationship O'Hare has calculated the heat of formation of  $\text{NS}$  and  $\text{NS}^+$  (Table 1). O'Hare's value of  $\Delta H_{\text{f}}(\text{NS}^+)$  was slightly higher than that obtained by Hobrock, Shankal and Kiser [55] (10.9 e.v.). He ascribed this difference between the two value to the incorrect assumptions made by Hobrock, et.al., in their calculations.

Ab initio calculations for  $\text{NS}$ ,  $\text{NS}^+$  and  $\text{NS}^-$  based on the Hartree-Fock Roothan approach by various workers [56-60] yielded the values of the ionization potential, and the dissociation energy in excellent agreement with the ones obtained by the experimental data.

It is interesting to note the  $D(NS)$  is less than that of  $D(NS^+)$ . Since  $NS^+$  has one less antibonding electron than in  $NS$ , the lower value of  $D(NS)$  is not unexpected. This type of lowering in the values of dissociation energies of the singly + vely charged species was also observed in other diatomic molecules which are isoelectronic with  $NS$  e.g.  $SiF$ ,  $PO$ ,  $CCl$  etc. The same trend is also observed in the values of this ionization potentials of  $NS$  and  $NS^+$ .

### The Electronic Structure of $NS$ and Related Compounds

It is interesting that in contrast to its homolog,  $NO$ , having the same number of valence electrons, thionitrosyl molecule has only a transient existence in the gaseous phase. In solid or liquid phase it polymerizes too readily into  $S_2N_2$ ,  $S_4N_4$  and  $(SN)_x$  and is therefore not possible to isolate it. Further,  $S_2N_2$ , unlike  $N_2O_2$ , a colourless crystalline solid, is highly explosive if not pure and undergoes polymerization into  $(NS)_x$  via  $S_4N_4$ . The geometries of these molecules are summerized in Table II. Several attempts [24-26, 64, 70-72] have been made to understand the electronic factors which lead to such a behaviour of  $NS$ .

Monomer NS is a paramagnetic molecule having a doublet  $^2\pi_{1/2}$  ground state with the lowest excited state,  $^2\pi_{3/2}$  lying  $223 \text{ cm}^{-1}$  above the ground level. A calculated electronic configuration of the ground state of NS is  $16^2 26^2 36^2 1 \pi^4 2 \pi^1$  where the valence electrons of only nitrogen and sulfur are considered. In the molecular orbital description of NO  $36^2$  and  $1 \pi^4$  levels are reversed, although the energy distance between them ( $36^2$  and  $1 \pi^4$ ) is small in both the molecules [73]. The calculations indicate that the sulfur d-orbital participation in the bond formation of NS is small ( $< 10\%$ ). With the result its effect on orbital energies is less than 0.3 e.v. (Fig. 1). However, for  $\text{S}_2\text{N}_2$  the extent of d orbital participation is relatively larger though still of minor importance. A charge distribution analysis in each of the MO as well as the total charges on sulfur and nitrogen in SN shows that the charges in the sulfur and nitrogen spheres are within a few percent of each other for all the molecular orbitals. Consequently, since the number of valence electrons on atomic sulfur is one more than that on atomic nitrogen, a net charge transfer (equivalent to approximately 0.5 electron) takes place from sulfur to nitrogen, which is in conformity with the electronegativity values of sulfur (2.5) and nitrogen (3.0). It will, thus, lead to a high value of the dipole moment of SN ( $\overset{+}{\text{S}} - \overset{-}{\text{N}}$ ) (Experimental value of dipole moment of SN is 1.86 D [68]). The situation in NO is just opposite

to that of  $\text{NS}$ . The charge distribution in nitrogen and oxygen spheres in  $\text{NO}$  is less even and the calculated value of charge transfer is approximately equivalent to 0.32 electrons.

This will result in a lower value of dipole moment of  $\delta^- \delta^+$   
 $\text{N} - \text{O}$  [the observed value of dipole moment of  $\text{NO}$  is 0.16D ].

The direction of dipole moment of  $\text{NO}$  is opposite to what one would expect on the basis of the electronegativity values of nitrogen (3.0) and oxygen (3.5). Hartree and Fock calculations [74,75] predicted that the dipole moment of  $\text{NO}$  will be small and of opposite sign to that predicted by methods which include correlation effects. Thus, the dipole moment of  $\text{NO}$  depends on the detailed behaviour of electrons rather than on the net electronic charge in a somewhat arbitrary volume around each nucleus.

The structure of  $\text{S}_2\text{N}_2$  (a planar square with alternating sulfur and nitrogen atoms and a bond distance of 1.65 Å) could be due to aligning of two  $\delta^+ \delta^-$   $\text{S} - \text{N}$  molecules in such a way that the two dipoles interact in an electrostatically favourable manner. This type of alignment will not be stable in case of  $\text{N}_2\text{O}_2$  which is actually a loosely bond trapezoidal arrangement of N and O with the shortest distance (2.18 Å) between two nitrogen atoms [76,84]. Despite the chemical binding in  $\text{S}_2\text{N}_2$  is more complex than the simple dipolar interaction, it seems that one of the main factors controlling

the relative geometry of  $S_2N_2$  and  $N_2O_2$  is electrostatic in nature. A charge distribution study in  $S_2N_2$  shows a net charge transfer of 0.48 electron per atom from S to N. A CNDO/2 localized MO study of  $S_2N_2$  leads to the conclusion that the molecule is formed from  $6^-$  electrons in three center bonds and four unshared pairs of electrons superimposed on a square planer ( $D_{2h}$  symmetry) bond structure [82-84]. The physical and chemical properties of disulfur dinitride have been subject of several studies [85-110].

A considerable amount of work (both theoretical and experimental) has been done on the structure of  $S_4N_4$  [111-131]. It is now generally agreed that  $S_4N_4$  has a structure with four coplanar nitrogen atoms and four sulfur atoms alternatively above and below the plane. However, the question of the electronic structure and the nature of bonding are still open. Various workers [118-121] have found a different net charge transfer from sulfur to nitrogen (0.328-0.56 electron). All the methods indicate substantial S-S orbital overlap and little or practically no N-N bonding interaction. All the studies of the sulfur nitrides relating to their structure lead to polar bonds in them. It, further, appears that in the polymerization process of SN to  $(SN)_x$  bond dipolar forces determine the geometry of the dimer,  $S_2N_2$  which is then

thermally excited to a low lying high spin state and the chain propagation takes place as a result of interaction of these excited species followed by their rearrangements to yield  $(\text{SN})_x$  structure.

(d) Spectroscopic Properties of NS

A large number of papers related to the microwave spectra of NS have appeared in the literature [132-165]. In 1932 Fowler and Bakker [21] have discovered the  $A_x$  and  $C_x$  systems of NS radical in the light emitted by an electric discharge through a mixture of nitrogen and sulfur vapour. Subsequently there have been a number of other investigations in the course of which the excited valence states  $B^2\pi$ ,  $A^2\Delta$ ,  $G^2\Sigma^-$ ,  $H^2\pi$  and  $I^2\Sigma^+$  have been identified besides the ground  $X^2\pi$ . Rosen has compiled

---

\* The dipole moments of  $\text{NS}^-$  and  $\text{NS}^+$  are 0.09D [33] and 3.893 D respectively. The change of dipole moments of NS from 1.84D (theoretical 1.73D [68]) to 0.09D for  $\text{NS}^-$  is consistent with the addition of one electron primarily at the sulfur end of the molecule.

all the spectroscopic data of NS molecule upto 1969 in the book "Spectroscopic Data relative to diatomic molecules." Furthermore a low lying  $2 \Sigma^+$  has also being reported which has been named as  $\beta$  state. The observed states, X, B, A, G, H and I are associated respectively with the theoretical states  $2\pi(1)$ ,  $2\pi(2)$ ,  $2\Delta(1)$ ,  $2\Sigma^-(1)$ ,  $2\pi(3)$  and  $2\Sigma^+(2)$ .

The electronic emission spectrum of NS has been discussed by Joshi, et.al. [22] . The electronic states of NO and NS are shown in Fig. II.

The i.r. spectra of NS radical showed a very weak band at  $1225 \text{ cm}^{-1}$  [30] (the vibration frequency of the gaseous NS molecule [70,166,167] is given as  $1204.1 \text{ cm}^{-1}$ ).

Very little information is available in the literature concerning the ionic state of NS, the only ionic state to be even partially characterized is the ground state,  $\Sigma^+$ . The band associated with the first ionization potential of the NS molecule corresponding to the ionization  $\text{NS}^+(\text{X}^1\Sigma^+) \longrightarrow \text{NS}(\text{X}^2\pi)$  has been observed by photoelectron spectroscopy using HeI and NeI photon sources [160]. The vertical ionization of this band has been measured as  $8.87 \pm 0.01 \text{ e.v.}$  and the position of three vibration components associated with this band leads to  $W_e = (1415 \pm 20) \text{ cm}^{-1}$  and  $W_0 = (15 \pm 10) \text{ cm}^{-1}$  for the  $\text{NS}^+(\text{X}^1\Sigma^+)$  state). The



increased value of  $W_e$  compared with the neutral molecule value of  $1219\text{ cm}^{-1}$  [72] is, as expected, consistent with removal of an electron from an antibonding orbital and is directly analogous to the corresponding increase in vibrational frequency on going from  $\text{NO}(X^2\Pi)$  to  $\text{NO}^+(X^1\Sigma^+)$

The potential energy diagram (Fig. III) has been constructed by using all the  $\text{NS}^+(^1\Sigma^+)$  parameters. An average of the parameters of the  $X^2\Pi_{1/2}$  and  $X^2\Pi_{3/2}$  states has been taken [72] in drawing this diagram. Also values of the dissociation energy and first ionization potential of  $\text{S}(^3\text{P})$  have been taken as  $(4.87 \pm 0.25)\text{ eV}$  [38, 64] and  $10.357\text{ eV}$  [63] respectively.

#### (e) Electron Paramagnetic Resonance

Carrington, et. al., first observed the electron paramagnetic resonance spectrum of NS obtained by the action of nitrogen atoms with hydrogen sulphide in a spectrophotometer cavity, [169-171] and by reacting N-atoms with  $\text{S}_2\text{Cl}_2$  [66]. Uehare and Merino [172] have detected an intense electron paramagnetic resonance spectrum of NS in the gas phase by passing a mixture of  $\text{N}_2$  and  $\text{SCl}_2$  through a microwave discharge cavity and then through a spectrometer cavity. The electron paramagnetic resonance spectrum of NS

consists of three triplets and the position of the centre of the spectrum indicates a g factor close to 0.8 (the value expected for a molecule in a  $2\pi_{3/2}$  state in the lowest rotational level with  $J=3/2$ ). The analyses of ultraviolet spectrum of NS shows that the ground state is actually  $2\pi_{1/2}$  state. The electron spin g factor  $g_e$  of NS lies probably between the free spin g factor and the electron spin g factors for  $O_2$  [173] or for SO [174]. (Fig.IV) NS radical would be expected to have an electronic structure similar to that of NO and molecular parameters for the two species are compared in Table III.

Table II. Geometries of SN, S<sub>2</sub>N<sub>2</sub> and S<sub>4</sub>N<sub>4</sub>

		r <sub>SN</sub>	SNS	NSN
SN	C <sub>∞v</sub>	1.495	-	-
S <sub>2</sub> N <sub>2</sub>	D <sub>2h</sub>	1.620	95	85
S <sub>4</sub> N <sub>4</sub>	D <sub>2d</sub>	1.616	113	105

Table III. Molecular Parameters for NS and NO

	NS	NO
Rotational constant B <sub>0</sub>	0.7724 cm <sup>-1</sup>	1.6957 cm <sup>-1</sup>
Fine structure constant A	223.0(3) cm <sup>-1</sup>	123.160±0.02 cm <sup>-1</sup>
Hyperfine constant h	56.8±0.5 MHz	75.81 ±0.24 MHz
Quadrupole coupling constant Q	3.1±0.5 MHz	1.8 ±0.3 MHz
Dipolemoment u	1.732 D	0.158±0.005 D




Fig.1 Orbital energies for the molecular orbitals of NS.

$\uparrow \rightarrow$  d-orbital contribution




Fig. II Electronic states for NO and NS molecules.




Fig. III Potential energy diagram










Fig. IV EPR spectrum of NS.

### Ionic Compounds

In 1971 Glemser and Koch [34,35] for the first time obtained the ionic compound of  $\text{NS}^+$  by the reaction of NSF with  $\text{AsF}_5$ . A number of other methods for the syntheses of such salt like compounds have since then been developed. The best results were achieved by reacting NSF with the Lewis acids  $\text{MF}_5$  [M=As or Sb] or  $\text{BF}_3$  in the gaseous phase at reduced pressure [36,37,175]. For preparative purposes these salts could best be synthesized by the reaction of NSF with  $\text{MF}_5$  in liquid  $\text{SO}_2$ . A yield upto 90-100% can be attained by this method.

The thionitrosyl compounds,  $[\text{NS}^+]$ ,  $[\text{AsF}_6]^-$ ,  $[\text{NS}^+[\text{SbF}_6]^-]$  or  $\text{NS}^+\text{BF}_4^-$ , were also obtained by the action of tri or tetramer of NSF with an excess of  $\text{AsF}_5$ ,  $\text{SbF}_5$  or  $\text{BF}_3$  respectively [176].





Purification of  $\text{NS}^+\text{AsF}_6^-$  can be affected by subliming the crude product at  $40^\circ\text{C}$  under reduced pressure whereby a stable colourless waxy solid which can be kept without decomposition for years in dry glass vessel, was obtained. The purification of  $\text{NS}^+\text{SbF}_6^-$  was accomplished by heating it at  $100^\circ\text{C}$  and the residue, thus left, is crystallized either from liquid  $\text{SO}_2$  or from  $\text{SO}_2\text{ClF}$ . Tetrafluoro derivative is not stable even at low temperatures and it could, therefore, not be purified by recrystallization.

The ionic nature of these compounds was confirmed by the following experimental techniques.

(a) Raman spectra : The Raman spectra of all these salts showed one band in the region  $1435\text{ cm}^{-1}$  which was assigned to  $\gamma_{(\text{NS})}$ . This band appeared in the spectrum of NSF at a lower frequency ( $1372\text{ cm}^{-1}$ ). From these band positions the valence field force constants\*\* of the NS and its bond order were calculated. The results indicate a considerable strengthening of NS bond in  $\text{NS}^+$  salt like compounds, which further confirms the ionic nature of the salts. The frequencies of some of the compounds are shown in Table IV.

---

\*\*  $f(\text{NS}^+)$  in  $\text{NS}^+\text{AsF}_6^- = 11.85\text{ Dynes/A}^\circ$      $f(\text{NS})$  in NSF =  $10.71\text{ Dynes/A}^\circ$

Bond order,  $n(\text{NS}^+) = 2.6$

Bond order of NS;  $n_{\text{NS}} = 2.4$

(b) NMR Spectra: There appeared no signal of  $F^{19}$  between +350 to -440 ppm, characteristic to SF. It has therefore been assumed that there is no bond between sulfur and fluorine.

(c) Exchange Reactions: The lack of exchange of  $F^{19}$  between the anions of these compounds and  $(AsF_6)^-$  further confirm the ionic nature of the compounds.

Since attempts to obtain a single crystal of these compounds failed, no single crystal x-ray structural studies was thus far reported in the literature.


Thionitrosyl salts have been used to introduce thionitrosyl group into other molecules. Synthetic possibilities by these ionic compounds include reactions with anions, nucleophiles and with compounds having polar bonds. Thus,

(i) Reaction of thionitrosyl cation with anions

The reaction of  $NS^+AsF_6^-$  with  $NO^+(CF_3SO_3)^-$  in liquid sulfur dioxide yielded a colourless product,  $NS^+[CF_3SO_3]^-$  (m.p.  $120^\circ$ ) which was purified by sublimation ( $30^\circ C/0.01$  torr). It was extremely hygroscopic and sensitive to hydrolysis. The ease of sublimation was taken as an indication for its being monomer at least in the gaseous phase. The presence of a band in its Raman spectrum due to  $\nu_{NS^+}$  at  $1443\text{ cm}^{-1}$  is in support of its being an ionic compound.

(ii) Reaction of thionitrosyl cation with nucleophiles

Insertion of NS unit into  $S_4N_4$  gives yellow pentathiazyl salts in the yields of about 80%. Thus,




M=As (yellow solid m.p.  $267^\circ\text{C}$ ) or Sb (yellow solid, m.p.  $175^\circ\text{C}$ )


The formulation of the intermediate product in these reactions is quite arbitrary because both are simply addition of the nucleophile,  $S_4N_4$ , to sulfur of  $NS^+$  to form either linear cationic thiazenes or a cyclo addition product of  $NS^+$  to  $S_4N_4$ . When the same reaction was carried out in liquid  $SO_2$ , the reaction product was a black solid,  $(S_5N_5)^+AsF_6^-$  (dec  $110^\circ\text{C}$ ) together with other unidentified substances. The i.r. spectrum of  $S_5N_5^+AsF_6^-$  showed a strong band at  $700\text{ cm}^{-1}$  characteristic of  $AsF_6^-$  group and a broad band at  $850\text{ cm}^{-1}$  indicating its polymeric nature. The ratio of  $NS:AsF_6$  in the black compound can be varied by changing the stoichiometry of the reacting species. A drop in this ratio (increase in charges of the polymer) was associated with a shift of position of  $\nu_{NS}$  band towards higher wave numbers.

(iii) Reaction of the thionitrosyl cation with polar bonds

The reaction of  $S_2Cl_2$  with  $NS^+MF_6^-$  gave  $N(SCl_2)^+MF_6^-$  ( $M=As$  or  $Sb$ ) according to the following reaction :



The synthesis of the cation,  $[N(SCl_2)]^+$  [177] was first accomplished as tetrachloroborate from the reaction of  $NSF_3$  and  $BCl_3$ . It can also be obtained from  $SCl_2$ ,  $(NSCl)_3$  and  $MCl_3$  [178,179]. Thus,



$M = Al, Sb, As$

The compound  $[N(SCl)_2]^+[MF_6]^-$  which can be regarded as derivative of:  $ClSN = SCl_2$  or  $ClSN = SClF$ , decomposes on reacting with  $NOCl$ .

The reaction of  $NS^+MF_6^-$  with  $SCl_2$  can be generalized for sulfenyl chloride. Thus, the salts  $[CF_3SNSCl]^+[MF_6]^-$  and  $[(CF_3)_2C=N-SNSCl]^+MF_6^-$  were prepared by reacting  $NS^+MF_6^-$  with  $CF_3SCl$  and  $(CF_3)_2C=N-SCl$  respectively [180].

Table IV. Thionitrosyl Cations

	m.p. °C	Colour	Raman frequency $\nu_{\text{NS}}$ cm <sup>-1</sup>	Reference
$\text{NS}^+\text{AsF}_6^-$	-	Colourless	1437	34,35
$\text{NS}^+\text{SbF}_6^-$	-	Colourless	1448	175
$\text{NS}^+\text{BF}_4^-$	-	Colourless	-	36,37
$\text{NS}^+\text{CF}_3\text{SO}_3^-$	120°C	Colourless	1443	175
$\text{NS}^+\text{AlCl}_4^-$	-	-	-	181
$\text{NS}^+\text{FeCl}_4^-$	-	-	-	181
$\text{NS}^+\text{GaCl}_4^-$	-	-	-	181

### Covalent Compounds


The chemistry of the compounds containing nitrogen sulfur and halide (halide = Cl or F) is mainly concerned with the covalent compounds of thionitrosyl. There are three principal covalent compounds of NS viz. NSF, NSF<sub>3</sub> and NSCl which we shall discuss here:

#### Thionitrosyl Fluoride NSF

The literature methods of synthesizing this compound are



##### (a) Fluorination of compounds containing NS bonds

AgF<sub>2</sub> [36,182], COF<sub>3</sub> [36] and HgF<sub>2</sub> [183] can fluorinate S<sub>4</sub>N<sub>4</sub> producing NSF.






Other compounds used as fluorinating agents are SeF<sub>4</sub>, F<sub>2</sub> [184], SF<sub>4</sub> [184,185], IF<sub>5</sub> and SbF<sub>5</sub>. The latter two agents give first adducts, viz (NSF)<sub>4</sub>S<sub>4</sub>N<sub>4</sub> or S<sub>4</sub>N<sub>4</sub>(SbF<sub>5</sub>)<sub>4</sub> which gave NSF when they are thermally decomposed [186].

##### (b) By reacting SF<sub>4</sub> with NH<sub>3</sub>






(c) By reacting  $\text{NF}_3$  with sulfur or metal sulfide [188-190]





In the course of these reactions other side products difficult to separate were also obtained. Maximum yield was found to be 20-30%.

(d) By decomposition of compounds containing NSF group [36,191-192]




The last reaction is suitable for the preparation of NSF in the large quantities.

Thionitrosyl fluoride (m.p.  $-89^\circ\text{C}$ , b.p.  $0.4^\circ\text{C}$ ) is a pungent smelling and pale yellow liquid. It is hydrolysed with water to give  $\text{NH}_3$ , sulfite and fluoride ion probably via unstable intermediate, thionylimide [194] .



The hydrolysis of NSF with dilute NaOH yielded first a yellow precipitate which was dissolved rapidly. It can not be stored at room temperature in glass vessels owing to its decomposition into  $\text{N}_2$ ,  $\text{SOF}_2$ ,  $\text{SiF}_4$  and  $(\text{NSF})_3$ . In teflon or copper containers it is relatively stable in liquid state but ~~there~~ trimerization also occur slowly.





In gaseous state at reduced pressure  $\text{S}_4\text{N}_4$  or  $\text{S}_3\text{N}_2\text{F}_2$  are the products of decomposition [195,196] .



Other reactions of NSF are given below :




(The reaction can be used to prepare  $\text{NSF}_3$ )



Reactions (ii) and (iii) illustrates the transformation of NSF into  $\text{NSF}_2$ .








Reactions (v) and vi) are used in the preparation of ionic compounds of  $NS^+$  ( see section on Ionic compounds)

(vii) The thionitrosylfluoride (NSF) is decomposed by HF [199]



(viii) The thionitrosylfluoride lose  $F^-$  and goes over to thiazyl cation



(ix)  $\gamma$ -irradiation of NSF in  $SF_6$  give probably  $NSF_2$  and FNSF radicals [200] .

Kirchhoff and Wilson [201] determined the molecular parameters from IR, NMR, microwave studies and proposed structure I . It was further confirmed by LCAOMO ab initio studies [202] and CNDO/2 calculations [203] . These results do not agree with the structure II and III proposed by Glemser, et.al. [204] from their studies and supported by Rogowski [205] on the basis of electron diffraction studies. Duke [202] recently observed that NSF is more stable than SNF. The observed S-F distance is 1.585Å in  $SOCl_2$ , 1.53Å in sulfonyl fluoride [206] , 1.58Å in  $SF_6$ , 1.50Å in  $S_2F_{10}$  and N S distance 1.416Å in  $NSF_3$ . It is evident from these

results that S-F and N S distances in thionitrosylfluoride are significantly larger than the ordinary single and triple bonds.




Fig. V

The NS force constant  $f(\text{NS})$  calculated from microwave and IR data has a value  $10.7 \text{ mdyne/\AA}^\circ$  [207-209] from which a bond order of 2.4 was calculated [210]. For NSCl having similar structure,  $f(\text{NS})$  is found to be  $10.0 \text{ mdyne/\AA}^\circ$  and bond order, 2.2 [211]. Thus, NS bond order depends upon the electronegativity of the element attached to sulfur which therefore suggests d - p bonding between N and S. Since then structure IV was proposed based on the hypothesis of Linnett [212]. This structure is thought to be the most stable configuration of NSF in the ground state as the doublet quartet of electrons [213]. The 'dots' represent the electrons with spin quantum number of  $+1/2$  and the 'crosses', the electrons with spin quantum number of  $-1/2$ .

or vice versa. Both spin sets of nine electrons favour a bent configuration. The two sets of four electrons around each nucleus can be treated as being uncorrelated and spatially related to one another. The electrostatic repulsion and the Pauli-principal effect cancel one another as far as the spatial correlation of the two sets is concerned. The electrons are widely separated in structure IV than they are in other structures.


The enthalpy of formation  $\Delta H_f(\text{NSF})$  gas, 298-15K)  $41 \pm 2 \text{ kcal mole}^{-1}$  [213] is derived from  $\Delta H_f(\text{NSF}_3, \text{ gas, } 298.15^\circ\text{K})$  and the mass spectrometric appearance potentials [214]. The dissociation energy  $D_{\text{NS}}(\text{NSF}) = 71 \pm 5 \text{ kcal mole}^{-1}$  [213]; dipole moment  $1.902\text{D}$  [207]. NMR and electronic spectral studies of NSF [215] have also been carried out.

### Thionitrosyl Trifluoride $\text{NSF}_3$


$\text{NSF}_3$  is the first compound with N-S triple bond and in which nitrogen and fluorine, together share six covalent bonds with sulfur.

$\text{NSF}_3$  can be synthesized (a) by fluorination of N-S compounds containing sulfur in +4 or lower oxidation state and (b) by the reaction of  $\text{S}^{+6}\text{-F}$  compounds with excess of ammonia. It can thus be obtained,


(a<sub>1</sub>) by refluxing tetrasulfurtetranitride with silver difluoride in carbon tetrachloride [36,183,197]




(a<sub>2</sub>) by fluorination of thionitrosyl fluoride with the silver difluoride [36,183,197,216]




(a<sub>3</sub>) by the reaction of COFNSF<sub>2</sub> with AgF<sub>2</sub> [198] (32)



(b<sub>1</sub>) by passing ammonia into a suspension of sulfur and AgF<sub>2</sub> in CCl<sub>4</sub>



and (b<sub>2</sub>) by reducing S<sub>2</sub>F<sub>10</sub> with excess of ammonia [216,218]




The last method of synthesizing NSF<sub>3</sub> is in particular not very much employed because of high toxicity of S<sub>2</sub>F<sub>10</sub> and of the low yield.


Thionitrosyltrifluoride is a colourless, pungent smelling gas (m.p.  $-72.6^{\circ} \pm 0.5$ , b.p.  $-27.1^{\circ} \pm 0.1^{\circ}$ ). Its enthalpy of formation and dissociation energy  $D(N\equiv S)$  are  $85 \text{ kcal mole}^{-1}$  and  $93 \pm 2 \text{ kcal mole}^{-1}$  respectively [224]. It is thermally and chemically very stable because of which its use as dielectric has been suggested [219]. It decomposes only above  $500^{\circ}\text{C}$  in glass vessels to give sulfur,  $\text{SO}_2$ , nitrogen,  $\text{SiF}_4$  and metallic fluorides as decomposition products [36]. It is unaffected by metallic sodium upto  $300^{\circ}\text{C}$  and unlike NSF it is not hydrolysed by dilute mineral acids.

The important reactions of  $\text{NSF}_3$  are as follows:

- (a) It is hydrolysed by water or by a strong alkali. In this reaction proton migration is suggested if a sulfur oxygen double bond can form at the expense of a sulfur-nitrogen multiple bond [220].




(b) It forms adducts with Lewis acids, like  $\text{BF}_3$ ,  $\text{MF}_5$  ( $\text{M}=\text{As}, \text{Sb}$ )  
 $[35, 37, 221]$




In gaseous phase these adducts dissociate into their components. Their i.r. spectra show a shift in the position of  $\nu_{(\text{NS})}$  towards higher wave numbers by about  $200 \text{ cm}^{-1}$  which is due to the increase in the N-S bond order from 2.7 to 2.9.

(c) It reacts with  $\text{LiN}[\text{Si}(\text{CH}_3)_3]_2$  to yield  $(\text{CH}_3)_3\text{Si}-\text{N}=\text{SF}_2=\text{N}-\text{Si}(\text{CH}_3)_3$  [222] which further reacts with one more molecule of  $\text{LiN}[\text{Si}(\text{CH}_3)_3]_2$  resulting in the splitting of two fluorine atoms attached to sulfur and the formation of the aza analogue of  $\text{SO}_3$  [222].



(d) The fluorine atom of  $\text{NSF}_3$  can be exchanged with  $-\text{NR}_2$  or  $-\text{OR}$  group while retaining its  $\text{N}-\text{S}$  bond [223]




$\text{R} = \text{C}_2\text{H}_5$  [224], piperidine [223]




$\text{R} = \text{C}_6\text{H}_5$  [223]


(e) Boron trichloride reacts with  $\text{NSF}_3$  with the formation of an ionic compound,  $[\text{N}(\text{SCl})_2]^+ [\text{BCl}_4]^-$



Although this reaction is not well understood mechanistically, but possibly it proceeds via an adduct formation followed by  $\text{F}-\text{Cl}$  exchange on sulfur and reduction of  $\text{NSF}_3$  to  $\text{NSCl}$ . The following syntheses of this compound support the reaction path




(f) Coordinatively unsaturated transition metal complexes react as Lewis acids and add electrophillically [224,225]



(M = Mn, Re)

These reactions are better carried out in liquid SO<sub>2</sub>, and in presence of only a small excess of NSF<sub>3</sub> e.g [225]




The thionitrosyltrifluoride complex  $[\text{M(CO)}_5\text{NSF}_3]^+ [\text{AsF}_6]^-$  (M = Re) is a colourless crystalline solid. The IR data of thionitrosyltrifluoride and a few of its coordination compounds are given in Table V.



Table V


	NSF <sub>3</sub>	F <sub>5</sub> AsNSF <sub>3</sub>	F <sub>5</sub> SbNSF <sub>3</sub>	Re(CO) <sub>5</sub> NSF <sub>3</sub>	[AsF <sub>6</sub> ] <sup>+</sup>	[Mn(CO) <sub>5</sub> NSF <sub>3</sub> ] <sup>+</sup>	[C <sub>5</sub> H <sub>5</sub> Fe(CO) <sub>2</sub> NSF <sub>3</sub> ] <sup>+</sup>	[AsF <sub>6</sub> ] <sup>-</sup>
$\nu_{\text{NS}}(\text{cm}^{-1})$	1515m	1650	1650	1636	1637	1632		
$\nu_{\text{SF}}(\text{cm}^{-1})$	811	950	955	900	882	888		
	775	889	900	889	838	876		
				846		830		





The primary cation, thus, formed should be unstable under the reaction conditions and it therefore takes  $\text{F}^-$  ion from the anion to form neutral  $\text{F}_3\text{S}-\text{N}=\text{SF}_4$ . Disproportionation then leads to  $\text{SF}_5-\text{N}=\text{SF}_2$ .

(h) The nitrogen of the thionitrosyltrifluoride is protonated by  $\text{HCl}$  in the presence of  $\text{AgAsF}_6$  [228]



(i)  $\text{LiN}(\text{SiMe}_3)(\text{CMe}_3)$  reacts with  $\text{NSF}_3$  to give  $\text{S}(\text{NSiMe}_3)(\text{:NCMe}_3)_2$  and  $\text{S}(\text{:NCMe}_3)_3$  [229].

### Structure

The structure of  $\text{NSF}_3$  (Fig. VI) has been determined from the IR [209, 230], Raman [231], NMR [209] and microwave spectra [232].




Fig VI

The NMR absorption of  $\text{NSF}_3$  [209] lies in the same region as that in the spectrum of  $\text{SF}_6$  suggesting all the fluorine atoms to be equivalent and bound to the sulfur atom. The molecule has, therefore,  $C_{3v}$  symmetry. This symmetry is further confirmed by I.R. spectrum [209, 230] which is similar to  $\text{OPF}_3$  as regards to the number and the shape of the bands. Calculation of the S-N bond strength from the force constants [209, 230, 231]  $f_{\text{NS}}(\text{NSF}_3)$  (12.4 mdyne/ $\text{\AA}^0$ ) and  $f_{\text{SF}}$  (5.6 mdyne/ $\text{\AA}^0$ ) give the bond order of N S; 2.7. These results were confirmed by the microwave spectrum of  $\text{NSF}_3$  [232]. It has a dipole moment of 1.91 Debye. The very short S-N bond distance indicates considerable amount of  $p\pi(\text{N}) - d\pi(\text{S})$  overlap (bond order 2.7). The electronic spectra of  $\text{NSF}_3$  has been discussed [231].

### Thionitrosyl chloride

The preparation of thionitrosyl chloride can be effected by any one of the following methods.

(a) By the action of chlorine on gaseous thionitrosylfluoride




(b) By the action of  $\text{S}_2\text{Cl}_2$  with ammonium chloride

(c) By thermal fission of  $\text{S}_3\text{N}_2\text{Cl}_2$ .


(d) By the action of chlorine upon  $S_3N_2Cl_2$  233, 234 .





(e) By reacting  $S_2Cl_2$  vapour and active nitrogen 233




(f) By the thermal decomposition of trithiazyltrifluoride  
 $N_3S_3Cl_3$  235, 236, 237 .




(g) By chlorination of tetrasulfurtetranitride [238]



(h) By the reaction of thionylchloride with thionylimide [239]



(i) By the reaction of  $H_2NS$  with trithiazyltrichloride [240]





### Properties

Thionitrosyl chloride, a greenish yellow gas, readily polymerizes to form a stable trimer [236, 241, 242] .



It is moisture sensitive and readily hydrolysed by water to form HNSO,  $\text{NH}_4^+$  and  $\text{SO}_3^{--}$  [239]



The values of the enthalpy and entropy change in the formation of NSCl from trithiazyltrichloride are  $\Delta H = 21.9 \pm 3.0$  kcal/mole and  $\Delta S^\circ(\text{NSCl}) = 77.5 \pm 9.4$  cal/deg/mole respectively.

NSCl is known to react with  $\text{S}_2\text{Cl}_2$  [233] according to the equation




The absolute entropy of NSCl(g) has been calculated from spectroscopic data [243] as  $63.66 \text{ cal deg}^{-1} \text{ mole}^{-1}$ .

The pressure of NSCl vapour in equilibrium with solid  $\text{N}_3\text{S}_3\text{Cl}_3$  is measured in a static system at  $31^\circ$  to  $60^\circ\text{C}$  [244].

A plot of  $\log P_{\text{NSCl}}$  vs  $\frac{1}{T}$  gives a straight line (Fig.VII) through the points corresponds to the equation

$P_{\text{NSCl}} = 12.321 - \frac{3360}{T}$ . From this  $H_f(\text{NSCl}) = 46.2 \pm 1.5$  kcal/mole and  $S(\text{NSCl}) = 129.6 \pm 4.8 \text{ cal deg}^{-1} \text{ mole}^{-1}$  were calculated for the reactions






Fig. VII Plot of  $\log P_{\text{NSCI}}$  vs.  $1/T$ .

The i.r. spectrum of NSCl [36,245-247] shows absorption bands due to  $\nu_{(\text{NS})}$ ,  $\nu_{(\text{SCl})}$  and  $\delta_{(\text{NSCl})}$  at  $1320 \text{ cm}^{-1}$ ,  $412 \text{ cm}^{-1}$ ,  $269 \text{ cm}^{-1}$  respectively.

### Structure

NSCl should be non linear according to Walsh rule 248 with  $C_s$  point group. Accordingly its i.r or Raman spectrum shows three allowed bands. From the positions of these bands ( $1320$ ,  $412$  and  $269 \text{ cm}^{-1}$ ), force constant were calculated by neglecting the cross terms in the potential function (Table VI). Because of the very high force constant, ( $f_{\text{NS}}=10.02 \text{ mdyne/\AA}^0$ ) [249] and bond order ( $\text{NS}_1=2.3$ ) values structures (I) and valence bond structures II & III and not the SNCl are proposed for thionitrosylchloride [247] .




Fig. VII



Table VI. Vibrational frequencies (in  $\text{cm}^{-1}$ ) and force constants (m dyne/ $\text{\AA}$ ) for NSCl

Vibration	Frequency	Fundamentals	Force constants
$\nu_{\text{NS}}$	1333.0(R)		
	1323.8(Q)	1324.5	$10.04 \pm 0.07$ ( $F_{11}$ )
	1316.0(P)		
$\nu_{\text{SCl}}$	423.9(R)		
	414.4(Q)	414.8	$1.49 \pm 0.23$ ( $F_{22}$ )
	405.6(P)		
$\nu_{\text{NSCl}}$	281.5(R)		
	275.2(Q)	271.6	$0.83 \pm 0.02$ ( $F_{33}$ )
	261.7(P)		

These structure are similar to those of thionitrosyl-fluoride. The structures of thionitrosylhalides are different from those of nitrosyl halides, NOX where halide atom is attached to the nitrogen atom and not with the oxygen [250-256]. Microwave spectra [257] of the NSCl was observed by pumping the products of thermally decomposing  $N_3S_3Cl_3$  continuously through the cell. The rotation constants were obtained for  $N^{14}S^{32}Cl^{35}$ ,  $N^{14}S^{32}Cl^{37}$ , and  $N^{15}S^{32}Cl^{35}$  in ground vibrational state and from there the molecular dimensions were determined. These are summarized in Table VII.

Table VII. Molecular dimension of NSCl

Compound	$N^{14}S^{34}Cl^{35}$	$N^{14}S^{32}Cl^{37}$	$N^{15}S^{32}Cl^{35}$
$r_{NS}$	1.450	1.458	1.448
$r_{SCl}$	2.163	2.157	2.164
$\angle NSCl$	$117^{\circ}, 48'$	$118^{\circ}, 8'$	$117^{\circ}, 45'$

The S-Cl bond length of NSCl is longer than the sum of Pauling's covalent radii ( $1.04 + 0.99 = 2.03 \text{ \AA}^0$ ). This anomaly in the bond length seems consistent with the fact

that the quadrupole coupling constant of the Cl nucleus is smaller than the average value for many ordinary molecules. This indicates the ionic weak nature of S-Cl bond in the thionitrosyl molecule.

### Structural Correlations

Various attempts have been made to establish a relationship between the N-S bond distance ( $r_{NS}$ ) and the bond order,  $\eta_{NS}$  (Fig. X) in order to characterize the N-S bond in NSF, NSF<sub>3</sub> and NSCl. As a result of this a relationship between  $r_{NS}$  and  $f_{NS}$  has been found which takes the form  $f_{NS} = 145 \cdot r_{NS}^{-7.00}$ . However this linear dependence of  $r_{NS}$  and  $\eta_{NS}$  [14,124,258,259] shall be considered only as an approximation. The Table VIII shows physical properties and Table IX shows force constants, bond distances, bond orders bond energies and i.r. frequencies for NSF, NSF<sub>3</sub> and NSCl. The frequency observed for NSF(1372cm<sup>-1</sup>) suggests that its bond order is less than that in NSF<sub>3</sub>(1515cm<sup>-1</sup>) and greater than that in NSCl (1320 cm<sup>-1</sup>) (Fig. IX).

Table VIII. Physical Properties of NSF, NSF<sub>3</sub> and NSCl

Compound	NSF	NSF <sub>3</sub>	NSCl
Molecular weight	65.01	103.01	81.46
Colour	pale yellow	colourless	greenish yellow
M.P.	-89	-72.6±0.5	-
B.P.	+0.4±2.0	-27.1±0.1	-
Density	-	1.92	-
Molar heat of volatilisation (cal)	6052±30	5526±50	-
Enthalpy of formation      H <sub>f</sub>	41±2 kcal/mole	-85 kcal/mole	21.9±3.0 kcal/mole
Trouton's constant	22.1	22.5	-

Table IX. Force Constant, Bond distance, Bond order and I.R. frequencies for NSF, NSF<sub>3</sub> and NSCl

Compound	$r_{\text{NS}} (\text{\AA})$	$f_{\text{NS}}$ (mdyne/ $\text{\AA}$ )	$\eta_{\text{NS}}$	Bond energy $d_{\text{NS}}$	I.R. ** ( $\text{cm}^{-1}$ )
NSF	1.446	10.71	2.4	71	1372 640 366
NSF <sub>3</sub>	1.416	12.55	2.7	93	1515 ( $\nu_{\text{NS}}$ ) 811 775 $\nu_{\text{SF}}$
NSCl	1.46	10.02	2.2	-	1320 412 273

\*\* The frequencies are written in order  $\nu_{\text{NS}}$ ,  $\nu_{\text{SX}}$  and  $\delta_{\text{NSX}}$




Fig. IX Correlation of S-N stretching frequencies and bond lengths.





Fig. X Correlation of force constants and bond lengths.


# Metallic and Non Metallic Adducts and Compounds of Sulfur Nitride

Many reactions of  $S_4N_4$  or  $S_2N_2$  with the compounds of the transition metals, post transition metals and with Lewis acids give rise to compounds in which (i) there are S-N ligands or (ii) there are adduct molecule of  $S_4N_4$ . These have been well reviewed [1,6] and only the main results will be described here (Table X).

Two type of structures have been found for  $S_4N_4$  adducts complexes. X-ray structure determination of  $S_4N_4SbCl_5$  [260] and  $S_4N_4BF_3$  [261] show that in the solid state they are covalent with donor nitrogen and unidentate  $S_4N_4$ . Conductance measurements have established [262] that in solution some compounds are ionized. Several other structures are possible in the solid state and in solution, for instances, (I) [263], (II) [260] and especially for 'soft' Lewis acids, structures which include donor sulfur.



(I)



(II)

Fig. XI

Table X. Compounds of  $S_4N_4$  and  $S_2N_2$  and their melting point and colour

S.No.	Compound	Colour(m.p.)	References
1	$S_4N_4TiCl_4$	(RBr, 132)	264, 265, 266
2	$S_4N_42TiCl_4$	(RBr, - )	264 - 266
3	$S_4N_4TiBr_4$	(DBr, 138)	264, 267
4	$S_4N_4TiF_4$	(O, 120)	267
5	$S_4N_4TiI_4$	(B, 100)	267
6	$S_4N_4VCl_4$	(Br, 190)	263, 264
7	$S_4N_42FeCl_3$	(RBr, 80)	267
8	$Fe(NS)_4$	(B, - )	268, 271, 272
9	$Co(NS)_4$	(DV, - )	269, 271, 272
10	$Ni(NS)_4$	(DV, - )	270, 271, 272
11	$Cu(NS)_2$	(Br, - )	273
12	$(CuNS)_4$	(B, - )	273
13	$Cu(NS)_2Cl_2$	(B, - )	273
14	$S_4N_4CuCl$	(B, - )	274
15	$S_4N_4CuBr$	(B, - )	274
16	$S_4N_4CuCl_2$	(B, - )	274
17	$S_4N_4ZrCl_4$	(RO, 260)	264, 267
18	$S_4N_4NbCl_5$	GBr, 180)	264
19	$S_4N_4MoCl_5$	(Bl, - )	260, 275
20	$Ag(NS)_2$	(RBr, - )	273
21	$S_4N_4HfCl_4$	(R, 140)	267



22	$S_4N_4TaCl_5$	(R, 180)	264, 267
23	$S_4N_4WCl_4$	(Br, - )	274, 267
24	$S_4N_4WBr_4$	(DaBr, 251)	267
25	$S_4N_4WOCl_4$	(DaBr, - )	267
26	$Pt(N_4S_4)(PPh_3)_2$		276
27	$Pt(N_2S_2)(PPh_3)_2$		276
28	$Pt(NS)_4$	(DaBr, 144)	272
29	$Pt(N_2S_4)(PPh_3)_2$		276
30	$Hg(SN)_2$		277, 278
31	$Hg(SN)_2$		277, 278
32	$S_4N_4BF_3$	(DaR, 145-147)	279, 280
33	$S_4N_4BCl_3$	(OR, 138)	262, 264, 280
34	$S_4N_42BCl_3$	(OBr, - )	262, 264, 280
35	$S_4N_4BBr_3$		280
36	$S_4N_4AlCl_3$	(DaR, - )	281
37	$S_4N_4AlBr_3$	(O, 122°C)	281
38	$S_4N_42AlBr_3$	(OBr, 144)	265
39	$S_4N_42GaCl_3$	(DR, 100)	267
40	$S_4N_42InCl_3$	(RBr, 100)	267
41	$S_4N_4\frac{1}{2}SnCl_4$	(DR, 200-202)	264, 275
42	$S_4N_4\frac{1}{2}SnBr_4$	(DBr, 198-200)	282, 283
43	$S_4N_42SbF_5$		284, 285
44	$S_4N_44SbF_5$	(g, 145)	284, 285
45	$S_4N_4SbCl_5$	(DR, 160-162)	262, 264, 275
46	$S_4N_42SbCl_5$		262, 264, 275

47	$S_4N_4 2SbBr_3$	284
48	$S_4N_4 2SbI_3$	284
49	$S_4N_4 SO_3$	286, 287
50	$S_4N_4 2SO_3$	286, 287
51	$S_4N_4 4SO_3$	286, 287
52	$S_4N_4 SeCl_4$ (Y, 127-129)	262
53	$S_4N_4 Se_2Cl_2$ (Y, - )	265
54	$S_4N_4 SeCl_4 SO_3$	264
55	$S_4N_4 TeCl_4$ (DR, 140°C)	262
56	$S_4N_4 TeB_4$	288
57	$S_4N_4 TeCl_4 BCl_3$	264
58	$S_4N_4 TeCl_4 SbCl_5$	264
59	$S_4N_4 TeCl_4 SO_3$	264
60	$S_4N_4 SnCl_4 POCl_3$	289
61	$S_3N_2 CF_3 CON$ (Y, - )	290
62	$S_3N_4 AsPh_3$	291

within brackets the letters indicate the following colours and the numbers for melting point in °C.

B= black; Bl= blue; Br= brown; D= deep; Da= dark;  
Y= green; O= orange; R= red, V= violet; Y=yellow.

The chemistry of the  $N_2S_2^+[292]$ ,  $S_3N_3^-[293-296]$ ,  $S_4N_3^+[297-311]$ ,  $S_4N_4^-[123, 311-313]$ ,  $S_4N_5^-[295, 296, 314, 315]$ ,  $(S_5N_5)^+[311, 316, 317]$  and  $S_5N_6^-[315]$  have also been studied.

## Transition Metal thionitrosyl complexes


Recently, there has been a considerable upsurge in the study of transition metal thionitrosyl complexes. This is partially due to the increase in understanding of the way in which NS binds to a metal. Similar to nitrosyl complexes, there are three principal bonding modes in thionitrosyl complexes: (a) terminal, linear M-N-S groups (b) terminal, bent M-N-S groups and (c) bridging NS groups. In case (a) the thionitrosyl ligand can be represented as coordinated "NS<sup>+</sup>" and in (b), as coordinated "NS<sup>-</sup>".

It is convenient to classify MNS complexes by the number of d type electrons present in the complex. Thus, MNS complex with n number of d electrons of M plus the number of electrons in the  $\pi^*$  orbitals of NS (or more conveniently, n is the number of d electrons if the thionitrosyl is regarded as being coordinated as NS<sup>+</sup>) would be written as  $\{MNS\}^n$ . This classification scheme will be used throughout the remainder of this review.

### Seven Coordination

The compounds  $[Mo(NS)(S_2CNR_2)_3]$ , ( $R_2=Me_2, Et_2, (CH_2)_4$ ) have been shown to be seven coordinated [318,319]. These were prepared by treating dioxo complex,  $MoO_2(S_2CNR_2)_2$  with

trimethylsilylazide in acetonitrile under reflux. It seemed likely that this reaction proceeds via intermediate formation of a nitrido complex and subsequently the nitrido complex  $[\text{MoN}(\text{S}_2\text{CNR}_2)_3]$  reacts with elemental sulfur in refluxing acetonitrile to give the thionitrosyl complexes.



(yield could be improved by using propylene sulphide).

These complexes are yellow, stable in air, diamagnetic and nonconducting in 1,2-dichloroethane solution. The  $^1\text{H}$  nmr spectrum of  $[\text{Mo}(\text{NS})(\text{S}_2\text{CNMe}_2)_3]$  in nitrobenzene at room temperature shows a 1:2:3 triplet due to the dithiocarbonate methyl group. The structure of  $[\text{Mo}(\text{NS})(\text{S}_2\text{CNMe}_2)_3]$  has been described as pentagonal bipyramide with the thionitrosyl group in one of the apical positions. The M-N-S angle for this  $\text{MNS}^4$  complex is  $172.0(7)^\circ$  [320]. The structure is shown schematically in Fig. XII.




Fig XII

In the thionitrosyl complex the Mo-N bond length ( $1.74\text{\AA}^\circ$ ) is longer than in the nitride ( $1.64\text{\AA}^\circ$ ), as would be expected specially, if the thionitrosyl ligand behaves in a manner analogous to the  $\text{NO}^+$  ligand, as the almost linear Mo-N-S system implies. A comparison between thionitrosyl complex and the analogous molecule  $[\text{Mo}(\text{NO})(\text{S}_2\text{CNBu}_2)_3][321]$  is summarized in Table XI.

TABLE XI

Complex	Mo-N distance	M-N-X
$\text{MO}(\text{NS})(\text{S}_2\text{CNMe}_2)_3$	$1.73\text{\AA}^\circ$	$173.2^\circ$
$\text{MO}(\text{NO})(\text{S}_2\text{CNBu}_2^\text{n})_3$	$1.74\text{\AA}^\circ$	$172.0(7)$

The Mo-N bond distance and M-N-X angle are equal, within the limit of experimental error. Thus it would seem that the NS and NO ligands are quite similar. The N-S distance in the free NS ligand [322] or the value calculated from the sums of covalent radii for an N=S double bond or even values found in a variety of compounds purporting to contain N=S partially multiple bonds [323] is smaller than the N-S distance ( $1.59\text{\AA}^\circ$ ) in the  $[\text{Mo}(\text{NS})(\text{S}_2\text{CNMe}_2)_3]$  which is consistent with a considerable degree of M-NS  $\pi$  back bonding.


The thionitrosyl sulfur in these complexes can be alkylated with triphenylmethyl tetrafluoroborate to give  $[\text{Mo}(\text{NSCP}_3)(\text{S}_2\text{CNR}_2)_3] \text{BF}_4$  [319, 325] .

### Six Coordination

All six coordinate complexes so far prepared are mononitrosyls.

### Ruthenium Complexes

Only two thionitrosyl complexes of ruthenium, (one with triphenylphosphine and one with triphenylarsine), are so far known which were prepared by the reaction of  $\text{RuCl}_3 \cdot \text{XH}_2\text{O}$  with  $(\text{NSCl})_3$  in THF or in  $\text{CCl}_4\text{-CHCl}_3$  mixture in the presence of triphenylphosphine and triphenylarsine 325 . (In THF or in  $\text{CHCl}_3\text{-CCl}_4$  mixture, the trithiazyltrichloride probably act as a solvated monomer) .




The i.r. spectra showed a strong sharp absorption bands in the  $1290\text{-}1295 \text{ cm}^{-1}$  region assigned to  $\nu_{\text{NS}}$ . The complexes are air stable, non conducting in nitrobenzene and are

diamagnetic. The structure of  $\text{Ru}(\text{NS})\text{Cl}_3(\text{AsPh}_3)_2$  has been determined by X-ray crystallography [326]. Full details of structure have not yet been published. The Ru-N-S angle is  $179^\circ$  suggesting that the NS ligand is functioning as a three electron donor. Thus, these six coordinate  $(\text{RuNS})^6$  complexes have been regarded as complexes between  $\text{Ru}(\text{II})$  and  $\text{NS}^+$ . The metal ligand bond was considered, by analogy with the  $\text{NO}^+$ , to be formed by donation of electrons from nitrogen to the metal and back donation from the filled metal d-orbitals to  $\pi^*(\text{NS})$  orbitals.


### Rhodium Complexes

Six coordinate complexes of the type  $\text{Rh}(\text{CO})(\text{NS})\text{Cl}_2\text{L}_2$  ( $\text{L}=\text{PPh}_3$  or  $\text{AsPh}_3$ ) have recently been reported [327,328]. Trithiatrichloride reacts with  $\text{trans Rh}(\text{CO})\text{ClL}_2$  in tetrahydrofuran to produce the diamagnetic, nonelectrolyte,  $\text{Rh}(\text{CO})(\text{NS})\text{Cl}_2\text{L}_2$  complexes.




The reaction of  $\text{RhH}(\text{CO})(\text{PPh}_3)_3$  with  $(\text{NSCl})_3$  in carbon tetrachloride-chloroform mixture gives a deep green chloro-bridged complex,  $[\text{Rh}(\text{CO})(\text{NS})\text{Cl}_2(\text{PPh}_3)]_2$  which on reaction with excess triphenylphosphine or triphenylarsine in

dichloromethane yields yellowish orange complexes  $[\text{Rh}(\text{CO})(\text{NS})\text{Cl}_2\text{L}_2] \cdot 0.5\text{CH}_2\text{Cl}_2$   $\text{L}=\text{PPh}_3$  or  $\text{AsPh}_3$  as a result of cleavage of chloro bridge. Their infrared spectra showed absorption bands in the range  $2100\text{--}2110\text{ cm}^{-1}$  ( $\nu_{\text{CO}}$ ),  $1118\text{--}1120\text{ cm}^{-1}$  ( $\nu_{\text{NS}}$ ) and  $330\text{ cm}^{-1}$  (terminal  $\nu_{\text{Rh-Cl}}$ ). In the spectrum of the green complex a band appeared at  $260\text{ cm}^{-1}$  which is assigned to bridged  $\nu_{\text{Rh-Cl}}$ . The diamagnetism of complexes indicate the oxidation state of rhodium either +1 or +3. The high  $\nu_{\text{CO}}$  frequency ( $\nu_{\text{CO}}$ ,  $2100\text{ cm}^{-1}$ ) in these complexes as compared to that in original compound ( $\nu_{\text{CO}}$ ,  $1960\text{ cm}^{-1}$ ) suggested +3 oxidation state of the rhodium. The shifting of the  $\nu_{\text{CO}}$  band towards higher wave numbers may possibly be due to the strong  $\sigma$ -bonding trans effect of the trans  $\text{NS}^-$  group. Thus, these six coordinate, bent,  $(\text{RhNS})^8$  complexes have been regarded as complexes between  $\text{Rh(III)}$  and  $\text{NS}^-$ . The probable structures of  $\text{Rh}(\text{CO})(\text{NS})\text{Cl}_2(\text{PPh}_3)_2$  (I) and  $\text{Rh}(\text{CO})(\text{NS})\text{Cl}_2\text{L}_2$  (II) are



(I)



(II)


Fig. XIII






The reaction of compounds  $\text{Rh}(\text{CO})(\text{NS})\text{Cl}_2\text{L}_2$  with  $\text{NOCl}$  in  $\text{CH}_2\text{Cl}_2$  afforded corresponding nitrosyl complexes  $\text{Rh}(\text{CO})\text{Cl}_2\text{L}_2$   
 $(\text{L} = \text{PPh}_3 \text{ or } \text{AsPh}_3)$

### Rhenium Complexes



Reactions of  $\text{S}_2\text{Cl}_2$  with  $[\text{ReCl}_2(\text{PRPh}_2)_2\text{N}](\text{R}=\text{Ph or Pr}^n)$  give thionitrosyl complexes,  $\text{ReCl}_3(\text{NS})(\text{PRPh}_2)_2$ . The complexes  $[\text{ReX}_2\text{N}(\text{PR}_3)_3]$ ,  $(\text{PR}_3 = \text{PMe}_2\text{Ph, PEt}_2\text{Ph or PMePh}_2; \text{X}=\text{Cl or Br})$  react with half an equivalent of  $\text{S}_2\text{Cl}_2$  to give the thionitrosyl  $[\text{ReClX}(\text{NS})(\text{PR}_3)_3]$ . Reaction of nitrido complex  $[\text{ReCl}_2\text{N}(\text{PR}_3)_3]$  with an excess of  $\text{S}_2\text{Cl}_2$  or  $\text{SCl}_2$  gave the derivatives  $[\text{ReCl}_3(\text{NS})(\text{PR}_3)_2]$ . The formation of thionitrosyl from nitrido and  $\text{S}_2\text{Cl}_2$  is possibly according to the scheme given in equation (64) [319, 329].



$[\text{Re}(\text{NS})\text{Cl}_2(\text{PMe}_2\text{Ph})_3]$  reacts with  $\text{K}(\text{CNS})$  or  $\text{Na}(\text{S}_2\text{CNMe}_2)$ .



Reaction of  $S_2Cl_2$  with  $[ReCl(N)(dPPE)_2]Cl$  ( $dPPE = Ph_2PCH_2CH_2PPh_2$ ) yields  $[Re(NS)Cl(dPPE)_2]Cl$  which reacts with  $FeCl_3$  or  $NH_4PF_6$  to give  $FeCl_4^-$  or  $PF_6^-$  salts according to equations




The salt  $[Re(NS)Cl(dPPE)_2](S_2CNEt_2)$  is prepared by the reaction of  $[ReCl(N)(dPPE)_2]Cl$  and  $Na(S_2CNEt_2)$ .

$Re(NS)Cl(dPPE)_2L$  ( $L = Cl, FeCl_4$  or  $PF_6$ ) are yellow and have conductivities in nitrobenzene typical of 1:1 electrolytes. In the infrared, a single strong peak observed in the region  $1167-1185\text{ cm}^{-1}$  confirms the presence of the  $NS^+$  group in these compounds.

The hydrogen -1 n.m.r. spectrum of  $[ReNSCl_2(PMe_2Ph)_3]$  shows two triplets and a doublet in the tertiary phosphine alkyl group region, indicating a meridional configuration for the phosphine ligands with the NS group trans to Cl. An x-ray crystal structure analysis of the analogous nitrosyl complex  $[Re(NO)Cl_2(PMe_2Ph)_3]$  shows the M-N-O system to be essentially linear with  $Re-N-O$   $178.8(1.4)^\circ$  [330]. It is


therefore extremely likely that the thionitrosyl analogues are isostructural with linear NS groups.



(II)

Fig. XIV

The H n.m.r. spectrum of  $[\text{Re}(\text{NS})\text{Cl}(\text{S}_2\text{CNMe}_2)(\text{PMe}_2\text{Ph})]$  shows a singlet in the dithiocarbamate methyl region and a pair of doublets in the tertiary phosphin alkyl group region, indicating configuration(III).





(III)

Fig. XV

The complexes,  $[\text{Re}(\text{NS})\text{Cl}_3(\text{PR}_3)_2]$  are paramagnetic with magnetic moments in the range 1.84-2.01 B.M. These values are close to that found for the analogous complex  $[\text{Re}(\text{NO})\text{Cl}_3(\text{PMe}_2\text{Ph})_2]$ .

### Osmium Complexes

Green complexes of osmium have been reported as a result of reactions of  $\text{OsX}_3\text{NL}_2$  ( $\text{L}=\text{AsPh}_3, \text{PMe}_2\text{Ph}$ ) or  $\frac{1}{2}$  bipy (2,2' bipyridyl) with half an equivalent of  $\text{S}_2\text{Cl}_2$ . Addition of pyridine ( $\text{P}_Y$ ) to a methylene chloride solution of the product formed by reaction of salt  $(\text{NBu}_4) [\text{OsCl}_4\text{N}]$  with  $\text{S}_2\text{Cl}_2$ , gives green  $[\text{Os}(\text{NS})\text{Cl}_3(\text{P}_Y)_2]$  [319]. The osmium thionitrosyl complexes,  $[\text{Os}(\text{NS})\text{Cl}_3\text{L}_2]$  ( $\text{L}=\text{PPh}_3$  or  $\text{AsPh}_3$ ) were prepared by the action of trithiazyltrichloride in THF with osmium trichloride in presence of triphenylphosphine or triphenylarsine respectively [325]



These complexes are diamagnetic and non electrolyte. In their infrared spectrum, a strong band is observed in the range  $1270\text{--}1282 \text{ cm}^{-1}$  indicating that the thionitrosyl group is coordinated as  $\text{NS}^+$ . The  $^1\text{H}$  n.m.r. spectrum of  $[\text{Os}(\text{NS})\text{Cl}_3(\text{PMe}_2\text{Ph})_2]$  shows a triplet due to the tertiaryphosphine methyl groups. This is consistent with the Fig. XVI with trans virtually coupled phosphorus ligands.

TABLE XII. Six Coordinate thionitrosyl complexes

Compound	Colour	N-S stretching frequencies cm <sup>-1</sup>	Refs.
$\text{Ru}(\text{NS})\text{Cl}_3(\text{PPh}_3)_2$	red brown	1290	325
$\text{Ru}(\text{NS})\text{Cl}_3(\text{AsPh}_3)_2$	red brown	1295	325, 326
$\text{Rh}(\text{CO})(\text{NS})\text{Cl}_2\text{PPh}_3_2$	green	1118	327
$\text{Rh}(\text{CO})(\text{NS})\text{Cl}_2(\text{PPh}_3)_2 \cdot 0.5\text{CH}_2\text{Cl}_2$	orange red	1120	327, 328
$\text{Rh}(\text{CO})(\text{NS})\text{Cl}_2(\text{AsPh}_3)_2 \cdot 0.5\text{CH}_2\text{Cl}_2$	orange red	1120	328
$\text{Re}(\text{NS})\text{Cl}_2(\text{PMe}_2\text{Ph})_3$	pink	1180	319
$\text{Re}(\text{NS})\text{Cl}_2(\text{PEt}_2\text{Ph})_3$	pink	1167	319
$\text{Re}(\text{NS})\text{Cl}_2(\text{PMePh}_3)_3$	pink	1172	319
$\text{Re}(\text{NS})\text{ClBr}(\text{PEt}_2\text{Ph})_3$	pink	1168	319
$\text{Re}(\text{NS})\text{Cl}(\text{SCN})(\text{PMe}_2\text{Ph})_3$	red pink	1177	319
$\text{Re}(\text{NS})\text{Cl}(\text{S}_2\text{CNMe}_2)(\text{PMe}_2\text{Ph})_2$	yellow brown	1150	319
$[\text{Re}(\text{NS})\text{Cl}(\text{dppe})_2]\text{Cl}$	yellow	1185	319
$[\text{Re}(\text{NS})\text{Cl}(\text{dppe})_2]\text{PF}_6$	yellow	1177	319
$[\text{Re}(\text{NS})\text{Cl}(\text{dppe})_2]\text{FeCl}_4$	yellow	1173	319
$[\text{Re}(\text{NS})\text{Cl}(\text{dppe})_2]\text{S}_2\text{CNEt}_2$	yellow	1183	319
$[\text{Re}(\text{NS})\text{Cl}_3(\text{PMe}_2\text{Ph})_2]$	purple	1228	319
$[\text{Re}(\text{NS})\text{Cl}_3(\text{PEt}_2\text{Ph})_2]$	purple	1230	319
$\text{Re}(\text{NS})\text{Cl}_3(\text{PMePh}_2)_2$	purple	1220	319
$\text{Re}(\text{NS})\text{Cl}_3(\text{PPh}_3)_2$	purple	1214	319

$\text{Re}(\text{NS})\text{Cl}_3(\text{PPr}^n\text{Ph}_2)_2$	purple	1226	319
$\text{Os}(\text{NS})\text{Cl}_3(\text{PPh}_3)_2$	green	1285	325
$\text{Os}(\text{NS})\text{Cl}_3(\text{AsPh}_3)_2$	green	1282	319, 325
$\text{Os}(\text{NS})\text{Cl}_3(\text{PMe}_2\text{Ph})_2$	green	1285	319
$\text{Os}(\text{NS})\text{Cl}_3(\text{bipy})$	green	1282	319
$\text{Os}(\text{NS})\text{Cl}_3(\text{P}_4)_2$	yellow green	1284	319
$\text{Os}(\text{NS})\text{ClBr}_2(\text{AsPh}_3)_2$	green	1270	319
$\text{Os}(\text{NS})\text{ClBr}_2(\text{bipy})$	green	1280	319
$\text{Ir}(\text{CO})(\text{NS})\text{Cl}_2(\text{PPh}_3)_2$	orange	1120	331

---

that the Vaska's complex in solution takes up oxygen from the atmosphere to form oxygen complex which on reaction with  $(\text{NSCl})_3$  forms thionitritoiridium complex.

### Five Coordination

These thionitrosyl complexes are amongst the very few examples of five coordination they are thionitrosyl derivatives of rhodium triphenylphosphine of general formula  $\text{Rh}(\text{NS})\text{ClX}(\text{PPh}_3)_2$  ( $\text{X} = \text{Cl}, \text{Br}$ ) [332]. They are prepared by the reaction of  $\text{RhX}(\text{PPh}_3)_3$  with trithiazylchloride in THF as red brown amorphous solid which are diamagnetic and nonconducting in nitrobenzene.  $\text{Rh}(\text{NS})\text{Cl}_2(\text{PPh}_3)_2$  is also prepared by action of  $\text{Rh}(\text{NS})(\text{PPh}_3)_3$  with chlorine gas [331]. The i.r. spectra of these complexes showed a absorption band in the region  $1117\text{--}1120\text{ cm}^{-1}$  which was assigned to  $\nu_{\text{NS}}$ . By analogy with nitrosyl complex  $\text{Rh}(\text{NO})\text{Cl}_2(\text{PPh}_3)_2$  the thionitrosyl complex  $\text{Rh}(\text{NS})\text{Cl}_2(\text{PPh}_3)_2$  may have the structure.





Fig. XVII

They are covalent in nature and soluble in benzene, chloroform and dichloromethane.

### Four Coordination

There are two compounds in this class, one each of chromium and rhodium. The action of trithiazyltrichloride on Na  $\left[\eta^5\text{-C}_5\text{H}_5\text{Cr(CO)}_3\right]$  in tetrahydrofuran gave di-carbonyl ( $\eta^5$  cyclopentadienyl) thionitrosylchromium(I)  $\eta^5\text{-C}_5\text{H}_5\text{Cr(CO)}_2\text{(NS)}$  as dark red violet, diamagnetic solid [333-335]



Its i.r. spectrum shows three strong bands attributable to terminal CO and NS groups.

The structure of  $\left[\eta^5\text{-(C}_5\text{H}_5\text{)Cr(CO)}_2\text{NS}\right]$  has been determined by x-ray crystallography [334,335] and is shown schematically in (Fig.XVI). Its molecular geometry is similar to that exhibited by molecules  $\left[\eta^5\text{-C}_5\text{H}_5\text{Mn(CO)}_3\right]$  [336] and  $\left[\eta^5\text{-C}_5\text{H}_5\text{Cr(CO)}_2\text{NO}\right]$  [338] and the Cr-C (cp), Cr-C (O) and C-O bond lengths are comparable to those found in other cyclopentadienyl chromium carbonyls [337]. The Cr-N-S angle is  $176.8(2)^\circ$  (Table XIII) suggesting that the thionitrosyl ligand coordinates essentially linearly to the metal via the nitrogen atom. Hence the thionitrosyl group can be considered to be bonded as  $\text{NS}^+$ , a mode of coordination that is directly analogous to metal-thiocarbonyl bonding



A comparison of the spectroscopic properties of the compounds  $(\eta^5\text{-C}_5\text{H}_5)\text{Cr}(\text{CO})\text{NX}(\text{X}=\text{O}, \text{S})$ , suggests that the NS ligand is more effective in removing electron density from the central atom than is the NO ligand.





Fig. XVIII

The reaction of  $(\eta^5\text{-C}_5\text{H}_5)\text{Cr}(\text{CO})_2(\text{NS})$  with  $\text{NOCl}$  and  $\text{NOPF}_6$  gave the nitrosyl complex  $(\eta^5\text{-C}_5\text{H}_5)\text{Cr}(\text{NO})_2\text{Cl}$  and a water sensitive, green black solid



The rhodium thionitrosyl  $\text{Rh}(\text{NS})(\text{PPh}_3)_3$  is prepared by action of trithiazyltrichloride in THF with hydrated rhodiumtrichloride in the presence of excess of triphenylphosphine and granulated zinc [331]





Table XIII. Four Coordinate Thionitrosyl Complexes

Complex	Colour	m.p. (°C)	I.R. frequencies		
			$\nu_{\text{CO}}(\text{cm}^{-1})$	$\nu_{\text{NS}}(\text{cm}^{-1})$	$\nu_{(\text{NO})}(\text{cm}^{-1})$
$(\eta^5\text{-C}_5\text{H}_5)\text{Cr}(\text{CO})_2(\text{NS})$	Red violet	68-69	2033, 1962	1180	-
$(\eta^5\text{-C}_5\text{H}_5)\text{Cr}(\text{CO})(\text{NO})(\text{NS}) \text{PF}_6$	Green black	-	2122	1243	1790
$\text{Rh}(\text{NS})(\text{PPh}_3)_3$	Pink	155	-	1120	-

TABLE XIV. Crystallographic Data on (  $\eta^5$ -C<sub>5</sub>H<sub>5</sub> )-Cr(CO)<sub>2</sub>NS

Bonds and angles	Values A or degrees
Cr-N	1.694(3)
Cr-C(O)	1.883(3)
C-O	1.131(3)
N-S	1.551(3)
$\angle$ Cr-N-S	176.8 (2)
$\angle$ Cr-C-O	178.1 (2)
$\angle$ C(O)-Cr-C(O)	92.4 (2)
$\angle$ C(O)-Cr-N	94.8 (1)

Reaction of  $\text{Rh}(\text{NS})(\text{PPh}_3)_3$  with chlorine gas afforded a reddish brown compound  $\text{Rh}(\text{NS})\text{Cl}_2(\text{PPh}_3)_2$




It is pink coloured, diamagnetic and nonconducting crystals. Its i.r. spectrum showed a strong absorption band at  $1120 \text{ cm}^{-1}$  which is assigned to  $\nu_{\text{NS}}$ .

#### Bridged thionitrosyl complex

The reaction of a carbontetrachloride solution of  $(\text{NSCl})_3$  with  $\text{Rh}(\text{NO})\text{Cl}_2(\text{PPh}_3)_2$  in  $\text{CHCl}_3$  under argon atmosphere produces a thionitrosyl bridged complex  $[\text{Rh}(\text{NS})\text{Cl}_2(\text{PPh}_3)]_2$  [338] which is air stable diamagnetic and nonelectrolyte in solution. Its i.r. spectra (Fig. XIX) did not show any band due to  $\nu_{(\text{NO})}$  at  $1630 \text{ cm}^{-1}$  but a new slightly broad band at  $840 \text{ cm}^{-1}$  appeared which was attributed to the bridging thionitrosyl groups. These results were interpreted as the replacement of NO group in the compound by the NS. The broadening of the  $840 \text{ cm}^{-1}$  band was suggested to be due to the two NS bands  $\nu_{\text{as}}(\text{NS})$  and  $\nu_{\text{sy}}(\text{NS})$  appearing close together. NS bridge in the compound were further


proved by reacting it with  $\text{PPh}_3$  and  $\text{AsPh}_3$  which resulted in the NS bridge cleavage.



(L =  $\text{PPh}_3$     brown m.p.  $145^\circ\text{C}$ )

(L =  $\text{AsPh}_3$     red brown m.p.  $160^\circ\text{C}$ )

The i.r. spectra of  $\text{Rh(NS)Cl}_2\text{(PPh}_3\text{)L}$  (L= $\text{PPh}_3$  or  $\text{AsPh}_3$ ) showed a characteristic band of terminal NS group at  $1120\text{ cm}^{-1}$  which further conforms with the fact that the band at  $840\text{ cm}^{-1}$  in complex  $[\text{Rh(NS)Cl}_2\text{(PPh}_3\text{)}]_2$  is a characteristic band of bridging thionitrosyl group.



A.  $\text{Rh}(\text{NO})\text{Cl}_2(\text{PPh}_3)_2$

B.  $[\text{Rh}(\text{NS})\text{Cl}_2(\text{PPh}_3)]_2$

C.  $\text{Rh}(\text{NS})\text{Cl}_2(\text{PPh}_3)_2$

D.  $\text{Rh}(\text{NS})\text{Cl}_2(\text{PPh}_3)(\text{AsPh}_3)$

References

- 1 H.G. Heal; Adv. Inorg. Chem. Radiochem. 15, (1972) 376.
- 2 M. Beeke-Goehring, Inorg. Chem. 1, (1959) 207.
- 3 M. Becke-Goehring, "Ergebnisse und Probleme der chemie der Schwefel stickstoffverbindungen", Akademie Verlag Berlin 1957.
- 4 M. Becke-Goehring, Adv. Inorg. Chem. Radiochem. 2, (1962) 159.
- 5 M. Becke-Goehring, Inorganic Macromol. Rev. 1, (1970) 17
- 6 M. Becke-Goehring and E. Fluck, Develop. Inorg. Nitrogen Chem. 1, (1966) 150.
- 7 H.W. Roesky, "Sulfur in Organic and Inorganic Chemistry", (A. Seening) Vol. 1 p.13 Dekker New York.
- 8 A.G. MacDiarmid, C.M. Mikulski, M.S. Saran; P.J. Russo M.J. Cohen; A.A. Bright; A.F. Gorito; A.J. Heeger, Adv. Chem. Ser. 150 (1976) 62.
- 9 D.K. Padma and A.R.V. Murthy, J. Sci Ind. Res. 35(5) (1976) 313.
- 10 A.J. Banister, 'Phosphorus Sulfur', 6(3) (1979) 421.
- 11 R. Appel and J. Kohnke, Method. Chim. 7, (1978) 743. Edited by Korte Friedhelm Academic New York N.Y.
- 12 I. Haidue Method. Chim. 7 (1978) 789, Edited by K. Friedhelm Academic New York N.Y.
- 13 H.G. Heal in 'Inorganic Sulfur Chemistry', (G. Nickless ed) p.459 Elsevier Amsterdam 1968.
- 14 O. Glemser, Angew. Chem. Int. Ed. Eng. 2 (1963) 530.
- 15 O. Glemser, Prep. Inorg. React. 1 (1964) 227.
- 16 O. Glemser and M. Field, Halogen Chem. 2 (1967) 1.
- 17 O. Glemser and U. Biermann; Nachr Akad. Wiss. Gottingen Math. Phys. Kl. 2 No.5 (1968) p.65.

References

- 1 H.G. Heal; Adv. Inorg. Chem. Radiochem. 15, (1972) 376.
- 2 M. Beeke-Goehring, Inorg. Chem. 1, (1959) 207.
- 3 M. Becke-Goehring, "Ergebnisse und Probleme der chemie der Schwefel stickstoffverbindungen", Akademie Verlag Berlin 1957.
- 4 M. Becke-Goehring, Adv. Inorg. Chem. Radiochem. 2, (1962) 159.
- 5 M. Becke-Goehring, Inorganic Macromol. Rev. 1, (1970) 17
- 6 M. Becke-Goehring and E. Fluck, Develop. Inorg. Nitrogen Chem. 1, (1966) 150.
- 7 H.W. Roesky, "Sulfur in Organic and Inorganic Chemistry", (A. Seening) Vol. 1 p.13 Dekker New York.
- 8 A.G. MacDiarmid, C.M. Mikulski, M.S. Saran; P.J. Russo M.J. Cohen; A.A. Bright; A.F. Gorito; A.J. Heeger, Adv. Chem. Ser. 150 (1976) 62.
- 9 D.K. Padma and A.R.V. Murthy, J. Sci Ind. Res. 35(5) (1976) 313.
- 10 A.J. Banister, 'Phosphorus Sulfur', 6(3) (1979) 421.
- 11 R. Appel and J. Kohnke, Method. Chim. 7, (1978) 743. Edited by Korte Friedhelm Academic New York N.Y.
- 12 I. Haidue Method. Chim. 7 (1978) 789, Edited by K. Friedhelm Academic New York N.Y.
- 13 H.G. Heal in 'Inorganic Sulfur Chemistry', (G. Nickless ed) p.459 Elsevier Amsterdam 1968.
- 14 O. Glemser, Angew. Chem. Int. Ed. Eng. 2 (1963) 530.
- 15 O. Glemser, Prep. Inorg. React. 1 (1964) 227.
- 16 O. Glemser and M. Field, Halogen Chem. 2 (1967) 1.
- 17 O. Glemser and U. Biermann; Nachr Akad. Wiss. Gottingen Math. Phys. Kl. 2 No.5 (1968) p.65.



- 18 O. Glemser, Endeavour 28 (1969) 86.
- 19 O. Glemser and R. Mews Adv. Inorg. Chem. Radiochem. 14 (1972) 14.
- 20 R. Mews, Adv. Inorg. Chem. Radiochem. 19 (1976) 185.
- 21 A. Fowler and C.J. Bakker, Proc. Roy. Soc. Ser. A 136 (1932) 28.
- 22 K.C. Joshi, Z. Phys. Chim (Frankfurt) 55 (1967) 173.
- 23 N.A. Narasimham and K.B. Subramanian J. Mol. Spectrosc 29 (1969) 294.
- 24 M. Peyron and L. Thanmy, J. Chim Phys. Chim. Biol. 64 (1967) 129.
- 25 J.A.S. Bett and C.A. Winkler, J. Phys. Chem. 68 (1964) 250.
- 26 P. Goudmand and O. Dessaux, J. Chim. Phys. 64 (1967) 135.
- 27 G. Ponnetier, P. Goudmand, O. Dessaux and N. Tavernier, C.R. Acad. Sci. 255 (1962) 91.
- 28 W.D. McGrath and T. Morrow, Nature 212 (1966) 5063.
- 29 B. Petropules; O. Dessaux; P. Goudmand, C.R. Acad. Sc. Ser. C. 270 (1970) 1223.
- 30 R.A. Teichman and R.E. Nixon; Inorg. Chem. 15 (1976) 1993.
- 31 R.D. Smith, J. Chem. Soc. Dalton Trans. (1979) 478.
- 32 K. Dressler, Helv. Phys. Acta. 28 (1955) 563.
- 33 P.A.G. O'Hare, J. Chem. Phys. 54 (1971) 4129.
- 34 O. Glemser and W. Koch, Angew. Chem. Int. Ed. Eng. 10 (1971) 127.
- 35 O. Glemser and W. Koch, Ann. Asoc. Guim Argen. 59 (1971) 143.
- 36 O. Glemser and H. Richert, Z. Anorg. Allgem. Chem. 307 (1961) 313.
- 37 O. Glemser; H. Richert and H. Haeseler, Angew. Chem. 71 (1959) 524.

- 38 P.B. Zeeman, Can. J. Phys. 29 (1951) 174.
- 39 P.A.G. O'Hare, J. Chem. Phys. 52 (1970) 2992.
- 40 Tate, Smith and Vaughan, Phys. Rev. 48 (1935) 525.
- 41 Hagstrum and Tate, Phys. Rev. 59 (1941) 59.
- 42 Angus and Leckie, Trans. Faraday Soc. 31 (1935) 958.
- 43 Londolt-Bornstein, "Tabellen" Band I Teil 3 (Atom-4 Molecular Physik), p.359 (1951).
- 44 L. Pauling, "The Nature of the Chemical Bond", Cornell University Press Ithaca N.Y. 1960 3rd Ed. p.78.
- 45 D.L. Hildenbrand and E. Murad, J. Chem. Phys. 51 (1969) 807.
- 46 A. Gaydon, "Dissociation and Spectra of Diatomic Molecules", Chapman & Hall 3rd Ed. (1968).
- 47 R.S. Berry, Chem. Rev. 69 (1969) 533.
- 48 A.C. Wehl and P. Maldonado Argonne National Laboratory (Private Communication) Hartree Fock Energy of NO = -129.2446.
- 49 E. Clementy A.D. Melean D.L. Raimondi and M. Yoshimine, Phys. Rev. 133 (1964) 1274.
- 50 J. Berkowitz, Argonne National Laboratory (Private Communication).
- 51 JANAF, Thermo chemical Tables (Dow Chemical Midland, Mich. (1963).
- 52 A.C. Wahl, P.J. Bertoneini, K. Kaiser and R.H. Land, USAEC Report No. ANL-7271 (1968).
- 53 G. Hertzberg, "Molecular Spectra and Molecular Structure", (D. Van Nortrand Co. New York 2nd Ed. (1950).
- 54 C.W. Beckett and E.C. Cassidy, Natt Bur. Std. U.S. Rept. (1969) 8628.
- 55 B.G. Hob-rock, R.C. Shenkel and K.W. Kiser, J. Phys. Chem. 67 (1963) 1684.
- 56 C.J. Roothaan, Rev. Modern Phys. 23 (1951) 69.

- 57 D.R. Salahub and R.P. Messmer, *J. Chem. Phys.* 64 (1976) 2039.
- 58 J.A. Jafri, M.D. Newton, T.A. Pakkanen and J.L. Witten, *J. Chem. Phys.* 66 (1977) 5167.
- 59 P.W. Deutsch and L.A. Curtiss, *Chem. Phys. Letters* 51 (1977) 125.
- 60 A. Karpfen P. Schuster J. Petkov and H. Lischka, *J. Chem. Phys.* 68 (1978) 3884.
- 61 JANAF Thermochemical Tables The Dow Chemical Company Milcland. Mich. (1965).
- 62 M. Icauss Natt, Bur. Btd (U.S.) Tech. Note 438 (1967).
- 63 C.E. Moore, "Ionization Potential and Ionization Limits", Derived from the analysis of optical spectra NSDRDS-NBS, 34 (1970).
- 64 Chi-Lian Chiu and S.J. Silvers, *J. Chem. Phys.* 63 (1975) 1095.
- 65 R. Byfleet, A. Carrington and D.K. Russell, *Mol. Phys.* 20 (1971) 271.
- 66 A. Carrington, B.J. Howard, D.H. Levy and J.C. Robertson, *Mol. Phys.* 15 (1968) 187.
- 67 T. Amano, S. Saito, Y. Monino, D.R. Johnson and F.X. Powell, *J. Mol. Spectrosc.* 30 (1969) 275.
- 68 P.A.G. O'Hare (Private Communication).
- 69 Y. Morino, Sagami Chemical Research Center Kahogawa Ven Japan (Private Communication).
- 70 R.F. Barrow, G. Drummoud and P.B. Zeeman, *Proc. Phys. Soc. A* 67 (1954) 365.
- 71 S.J. Silvers and C. Chiu, *J. Chem. Phys.* 61 (1974) 1475.
- 72 A. Jenouvrier and B. Pascot, *Can. J. Phys.* 51 (1973) 2143.
- 73 P. Siegbahn and H. Siegbahn (Unpublished).
- 74 R.M. Neumann, *Astrophys J.* 161 (1970) 779.
- 75 S. Green, *Chem. Phys. Lett.* 13 (1972) 552.

- 76 S. Green, *Chem. Phys. Lett.* 23 (1973) 115.
- 77 W.J. Dulmage, E.A. Meyers and W.N. Lipscomb, *Act. Cryst.* 6 (1953) 760.
- 78 W.N. Lipscomb, F.E. Wang, W.R. May and E.L. Lippert, *Acta Cryst.* 14 (1961) 1100.
- 79 C.E. Dinerman and G.E. Ewing, *J. Chem. Phys.* 54(1971)3660.
- 80 C.E. Dinerman and G.E. Ewing, *J. Chem. Phys.* 53(1970)626.
- 81 W.N. Lipscomb, *J. Chem. Phys.* 54 (1971) 3654.
- 82 J. Bragin and M.V. Evans, *J. Chem. Phys.* 51 (1969) 268.
- 83 J.R.W. Warm and D. Chapmann, *Spectrosc. Chem. Acta* 22 (1966) 1371.
- 84 A.G. MacDiarmid, C.M. Mikulski, P.J. Russo, M.S. Saran, A.F. Gorito and A.J. Heeger, *J. Chem. Soc. Chem. Comm.* (1975) 478.
- 85 C.M. Mikulski P.J. Russo M.S. Saran, A.G. MacDiramid, A.F. Garito and A.J. Helger, *J. Am. Chem. Soc.* 97 (1975) 6358.
- 86 U.V. Walatka, M.M. Labes and J.M. Perlsteen, *Phys. Rev. Lett.* 31 (1973) 1139.
- 87 R.L. Green, P.M. Grant and G.B. Street, *Phys. Rev. Lett.* 34 (1975) 89.
- 88 A.A. Bright M.J. Cohen A.F. Garito, A.J. Heeger, C.M. Mikulski, P.J. Russo and A.G. Mac Diarmid, *Phys. Rev. Lett.* 34 (1975) 206.
- 89 R.L. Green, G.B. Street and L.J. Setter, *Phys. Rev. Lett.* 34 (1975) 577.
- 90 V.T. Rajan and L.M. Falicov, *Phys. Rev. B* 12 (1975) 1240.
- 91 A. Zunger, *J. Chem. Phys.* 63 (1975) 4854.
- 92 L. Ley, *Phys. Rev. Lett.* 35 (1975) 1976.
- 93 W.E. Rudsc and P.M. Grant, *Phys. Rev. Lett.* 35(1975)1799.
- 94 P. Mengel, P.M. Grant, W.E. Rudge, B.H. Schechtman and D.W. Rice, *Phys. Rev. Lett.* 35 (1975) 1803.

- 95 R.L. Patton and W.L. Jolly, *Inorg. Chem.* 8 (1969) 1389.
- 96 R.L. Patton and W.L. Jolly, *Inorg. Chem.* 8 (1969) 1392.
- 97 R.L. Patton and K.N. Raymond, *Inorg. Chem.* 8 (1969) 2426.
- 98 R.N. Wiener, L.I. Rubin and S.N. Singh, U.S. At. Energy Comm. AFCRL-67-0512419 (1966).
- 99 D.R. Salahub and R.P. Messmer, *Phys. Rev. B* 14 (1976) 2592.
- 100 R.P. Messmer and D.R. Salahub, *Chem. Phys. Lett.* 41 (1976) 73.
- 101 M.P.S. Collins and B.J. Duke, *J. Chem. Soc. Chem. Comm.* (1975) 701.
- 102 F. Tarok, P. Pulay, T. Szondy and P. Nagy *Acta, Chim.* 80 (1974) 139.
- 103 A.J. Banister, M.T.P. *Int. Rev. Sci. Ser. 2, Inorg. Chem.* 3 (1975) 41.
- 104 W.L. Jolly, *ACS Chem. Ser.* 110 (1972) 92.
- 105 F. Cavallone and E. Clementi, *J. Chem. Phys.* 63 (1975) 4304.
- 106 M.J. Cohen, A.F. Garito, A.J. Heeger, A.G. MacDiarmid, C.M. Mikulski and M.S. Saran (Pre Print).
- 107 M. Kertesz, S. Seehai, A. Azman, D. Kocran and A.I. Kiss, *Chem. Phys. Lett.* 44 (1976) 53.
- 108 R.P. Messmer and D.R. Salahub, *Chem. Phys. Lett.* 41 (1976) 73.
- 109 R. Baughman, R.R. Chance and M.J. Cohen, *J. Chem. Phys.* 64 (1976) 1869.
- 110 M. Becke-Goehring and D. Voigt, *Z. Anorg. Allgen. Chem.* 285 (1956) 181.
- 111 E.R. Lippincott and M.C. Tobin, *J. Chem. Phys.* 21 (1953) 1559.
- 112 B.D. Sharma and J. Donhue, *Acta. Cryst.* 16 (1963) 891.
- 113 P.S. Braterman, *J. Chem. Soc.* (1965) 2297.

- 114 A.G. Turner and F.S. Mortimer, *Inorg. Chem.* 5 (1966) 906.
- 115 J.B. Mason, *J. Chem. Soc. A* (1969) 1567.
- 116 M.S. Gopinathan and M.A. Whitehead, *Can. J. Chem.* 53 (1975) 1343.
- 117 J.C. Slater in *advances in quantum chemistry* Vol.6 Edited by P.O. Lowdin Academic, New York, 1973, p.1
- 118 W.A. Goddard, "Modern Theoretical Chemistry", Edited by H.F. Schaefer (Plenum, New York) Vol. 1.
- 119 W.A. Goddard and R.C. Ladner, *J. Am. Chem. Soc.* 93 (1971) 6750.
- 120 P.J. Hay, W.J. Hunt and W.A. Goddard III, *J. Chem. Phys.* 57 (1972) 738.
- 121 W.J. Hunt, P.J. Hay and W.A. Goddard III, *J. Chem. Phys.* 57 (1972) 738.
- 122 M.J. Buerger, *Amer. Mineral* 21 (1936) 575.
- 123 D. Chapman and A.G. Massey, *Trans. Faraday Soc.* 58 (1962) 1291.
- 124 D. Chapman and T.C. Waddington, *Trans. Faraday Soc.* 58 (1962) 1679.
- 125 D. Clark, *J. Chem. Soc.* (1952) 1615.
- 126 W.P. Griffith and K.J. Rutt, *J. Chem. Soc. A* (1968) 2331.
- 127 C.S. Lu and J. Donobue, *J. Amer. Chem. Soc.* 66 (1944) 818.
- 128 J. Mason, *J. Chem. Soc. A* (1969) 1567.
- 129 R.A. Meinzer, D.W. Pratt and R.J. Myers, *J. Amer. Chem. Soc.* 91 (1969) 6623.
- 130 M.T. Rogers and K.J. Gross, *J. Am. Chem. Soc.* 74 (1952) 5294.
- 131 G.F.H. Smith, *Mineral Mag.* 16 (1913) 97.
- 132 N.A. Narasimham and K. Srikameswaran, *Proc. Indian. Acad. Sci. Sect. A* 56 (1962) 316.
- 133 N.A. Narasimhan and K. Srikameswaran *Nature*, 197 (1963) 370.

- 134 M.M. Patel, Z. Physik 173 (1963) 347.
- 135 G. Punnettier, P. Goudmand, O. Dessaux and N. Tauernier, J. Chim. Phys. 61 (1964) 395.
- 136 N.A. Narasimhan and K. Srikameswaran, Proc. Indian. Act. Sci. Sec. A59 (1964) 227.
- 137 J.J. Smith and B. Beyer, J. Mol. Spectra 14 (1964) 160.
- 138 J.A.S. Bett and C.A. Winkler, J. Phy. Chem. 68 (1964) 2735.
- 139 G. Pannetier, O. Dessaux, I. Arcliti and P. Goudmand, Compt. Rend. 259 (1964) 2198.
- 140 R.D. Young, R.L. Sharpless and R. Stringham, J. Chem. Phys. 40 (1964) 117.
- 141 A. Carrington and D.H. Levy, J. Chem. Phys, 44 (1966) 1298.
- 142 R.F. Barrow, G. Drummond and P.B. Zeeman, Proc. Phys. Soc. London A67 (1954) 35.
- 143 N.A. Narasimhan and T.K. Balasubramanian, J. Mole. Spectrosc. 40 (1971) 511.
- 144 K.C. Joshi, Z. Physik 191 (1966) 126.
- 145 G. Pannetier, P. Goudmand, O. Dessaux and I. Arditi, C.R. Congr. Nat. Soc. Sowautes Sect. Sci. 90(1) (1965) 117.
- 146 B. Pteropoulos, O. Desaux, D. Cherffioli and P. Goudmand, C.R. Acad. Sci. Paris, Ser. A, 13, 265 (1967) 355.
- 147 N.A. Narasimhan, T.K. Balasubramanian, Proc. Int. Conf. Spectrosc. Int. Bombay 1 (1967) 108.
- 148 B. Vival, O. Dessaux, J.P. Mattal and P. Gaudmand, C.R. Acad. Sci. Paris. Sci. C268 (1969) 2140.
- 149 F. Torok, P. Pulay, T. Szondy and P. Nagy, Acta. Chem. 80 (1974) 139.
- 150 N.A. Narasimhan and T.K. Balasubramanian, J. Mole. Spectrosc. 53 (1974) 128.
- 151 N.A. Narasimhan, K. Raughubeer and T.K. Balasubramanian, J. Mole. Spectrosc 54 (1975) 160.

- 152 F.J. Lovas, E.Jiemann, J. Phys. Chem. Ref. Data 3 (1974) 609.
- 153 H.R. Johnson, R.F. Beebe, and C. Sneden, Astrophys. J. Suppl. Ser. 29 (1975) 123.
- 154 Chi-Lian Chiu, Diss Abstr. Int. B36, (1975) 254.
- 155 C.A. Gottlieb, J.A. Ball, E.W. Gotlieb, C.J. Lada and H. Penfield, Astrophys. J. 200 (1975) 147- 149.
- 156 T.B.H. Kuiper, B. Zuckirman, R.K. Kakar and E.R. Kuiper, Astro. Phys. J. 200 (1975) 151- 153.
- 157 A. Jenouvrier and D. Daumont, J. Mol. Spectrosc. 61 (1976) 313.
- 158 Chi-Lian Chiu and S.J. Silvers, J. Mol. Spectrosc. 61 (1976) 316.
- 159 M. Vervloet and A. Jenouvrier, Can. J. Phys. 54(1976)1909.
- 160 J.M. Dyne, A. Morris and I.R. Trickle, J. Chem. Soc. Faraday Trans. 73 (1977) 147.
- 161 Ringers A. Douglas, U.S. NTIS AD Rep. 1977. AD-A037802 p.18, Avail NTIS Fron. Gov. Rep. Annouvle Index (U.S.) 77 (1977) 116.
- 162 D.A. Varshalovich and V.K. Khersonskii, Astron. Zh. 54 (1977) 915.
- 163 K. Raghuveer and N.A. Narasimham, J. Mol. Spectrosc. 70 (1978) 323.
- 164 Z. Iqbal and D.S. Dowas, Proc. Int. Conf. Lattics. Dyn. (1977) 582.
- 165 Murai Hisao, Yagi Toshikiko, Obi Kinichi, Tanako. Ikuzo Chem. Phys. Lett. 61 (1979) 531.
- 166 D. Chapman, R.J. Warm, A.G. Fitygerald and B.D. Fyoffe, Trans. Faraday. Soc. 60 (1964) 294.
- 167 G. Nicklss "Inorganic Sulfur Chemistry", Elesevier, New York (1968) p.
- 168 S.M. Duke, L. Golob, N. Sonathan and A. Morris, J. Chem. Soc. Faraday II. 71 (1975) 1026.



- 169 A. Carrington and D.H. Levy, J. Chem. Phys. 44 (1966) 1298.
- 170 A. Carrington and D.H. Levy, J. Phys. Chem. 71 (1967) 2.
- 171 A. Carrington, Proc. Roy. Soc. Ser.A 302 (1967) 2191.
- 172 H. Uehara and Y. Morino, Mol. Phys. 17 (1969) 239.
- 173 K.D. Bowers, R.A. Kamper and C.D. Lustig, Proc. R. Soc. A251 (1959) 565.
- 174 H. Uehors, Bull. Chem. Soc. Japan, 421 (1969) 886.
- 175 T. Mews, Angew. Chem. Int. Ed. Eng. 15 (1976) 691.
- 176 R. Mews, D.L. Wegner and O. Glemser, Z. Anorg. allgem. Chem. 412 (1975) 148.
- 177 O. Glemser, B. Krebs, J. Wegner and E. Kindler, Angew. Chem. Int. Ed. Eng. 8 (1969) 598.
- 178 O. Glemser and J. Wegner, Inorg. Nucl. Chem. Lett. 7 (1971) 613.
- 179 G.G. Alange, A.J. Banister and P.J. Dainty, Inorg. Nucl. Chem. Lett. 15 (1979) 175.
- 180 R. Mews, H. Steinbeisser and J. Varwig (Unpublished result).
- 181 A.J. Bonister and H.G. Clark, J. Chem. Soc. Dalton Trans. (1972) 2661.
- 182 O. Gleser, H. Richert and F. Rogowski, Naturewissenschaften 47 (1960) 94.
- 183 O. Glemser, H. Meyer and Hass, Chem. Ber. 97 (1964) 1704.
- 184 B. Cohen, T.R. Hooper, D. Heegil and R.D. Peacock, Nature, 207 (1965) 748.
- 185 O. Glemser and S. Austin (Unpublished observations 1961).
- 186 B. Cohen, T.R. Hooper and R.D. Reacock, J. Inorg. Nucl. Chem. 28 (1966) 919.
- 187 O. Glemser, E. Wysgomirski and H. Meyer (Unpublished observations 1964).

- 188 O. Glemser, U. Biermann, J. Knaak and A. Haas, Chem. Ber. 98 (1965) 446.
- 189 O. Glemser, J. Wegner and R. Mews, Chem. Ber. 100 (1967) 2474.
- 190 W.D. McGratch and T. Morrow, Nature (London) 212 (1966) 746.
- 191 A.F. Clifford, C.S. Kobayashi and J.H. Stanton, 148th Meet. Amer. Chem. Soc. Chicago Abstract K8 (1964).
- 192 J.K. Ruff, Inorg. Chem 5 (1966) 1787.
- 193 O. Glemser, R. Mews and H.W. Roesky, Chem. Ber. 102 (1969) 1523.
- 194 H.W. Roesky, O. Glemser and A. Hoff, Chem. Ber. 101 (1968) 1215.
- 195 O. Glemser, H. Schroder and E. Wyszomirski, Z. Anorg. Allgem. Chem. 298 (1959) 72.
- 196 O. Glemser and E. Wyszomirski, Angew. Chem. 69 (1957) 534.
- 197 O. Glemser and H. Schroder, Z. Anorg. Allgem. Chem. 284 (1956) 971.
- 198 O. Glemser and S. Gruhr (Unpublished observations (1966)).
- 199 R. Mews (Unpublished observations 1976).
- 200 A.R. Boate and K.F. Preston, Inorg. Chem. 17 (1978) 1669.
- 201 W.H. Kirchhoff and E.B. Wilson, J. Amer. Chem. Soc. 85 (1963) 1726.
- 202 M.P.S. Collins and B.J. Duke, J. Chem. Soc. Dalton Trans. (1978) 277.
- 203 D. Gonbeau and G. Pfister Gnillouzo, J. Chim. Phys. Chem. Biol. 73 (1976) 787.
- 204 O. Glemser, H. Richert and F. Rogowski, Naturewissenschaften, 47 (1960) 94.
- 205 F. Rogowski, Z. Phys. Chem. 27 (1961) 277.
- 206 O.R. Lide, D.E. Mann and R.M. Fristrom, J. Chem. Phys. 26 (1957) 734.

- 207 R.L. Cook and W.H. Kirchhoff, J. Chem. Phys. 47 (1967) 4521.
- 208 A.M. Mirri and A. Guarnieri, Spectrochim Acta Part A 23 (1967) 2159.
- 209 H. Richert and O. Glemser, Z. anorg. allgem chem. 307 (1961) 328.
- 210 H. Sierbert, Z. Anorg. allgem. Chem. 273 (1953) 170.
- 211 A. Muller, G. Nagarajan, O. Glemser, S.F. Cyrin and J. Wegner, Spectrochim Acta Part A 23 (1967) 2693.
- 212 J.W. Linnett, J. Am. Chem. Soc. 83 (1961) 2643.
- 213 P.A.G. O'Hore, W.N. Nubbard, O. Glemser, and J. Wegner, J. Chem. Thermodyn. 2 (1970) 71.
- 214 O. Glemser, A Muller, D. Bohler and B. Kerbs, Z. Anorg. Allgem. Chem. 357 (1968) 184.
- 215 T. Barrow, R.N. Dixon, O. Glemser and R. Mews, Trans. Faraday, Soc. 65 (1969 ) 2295.
- 216 A.F. Clifford and J.W. Thompson, Inorg. Chem. 5 (1966) 1424.
- 217 B. Cohen and A.G. Mac Diamid, Chem. Ind. (1962) 1866.
- 218 B. Cohen and A.G. MacDiamid, Inorg. Chem. 4 (1965) 1782.
- 219 R.E. Eibeck, U.S. Patent 3 (1968) 39009.
- 220 H.W. Roesky, O. Glemser, A. Holl and W. Koch, 3 (1967) 39.
- 221 A. Muller, O. Glemser and K. Sherf, Chem. Ber. 99 (1966) 3568.
- 222 O. Glemser and J. Wegener, Angew. Chem. Int. Ed. Eng. 9 (1970) 309.
- 223 O. Glemser, W. Koch, Z. Naturforsch B23 (1968) 745.
- 224 R. Mews and O. Glemser, Angew. Chem. Int. Ed. Eng. 44 (1975) 186.
- 225 C.S. Liu, R. Mews and O. Glemser (Unpublished observations)

- 226 A.F. Clifford and L.C. Duncon, *Inorg. Chem.* 5 (1966) 692.
- 227 A.F. Clifford and G.R. Zlilenger, *Inorg. Chem.* 8 (1969) 979.
- 228 R. Mews (Unpublished observations).
- 229 F.M. Tesky, R. Mews, O. Glemser, *Z. Anorg. Allgen. Chem.* 452 (1979) 103.
- 230 K. Rakaswamy, K. Sathiandan and F.F. Cleveland, *J. Mol. Spectrosc.* 9 (1962) 107.
- 231 A. Muller, A. Rnoff, B. Krebs, O. Glemser and W. Koch, *Spectrosc. Chem. Acta Part A25* (1969) 199.
- 232 W.H. Kirehoff and E.B. Wilson Jr., *J. Am. Chem. Soc.* 84 (1962) 334.
- 233 K.D. Maguire, J.J. Smith and W.L. Jolly, *Chem. Ind. (London)* 39 (1963) 1589.
- 234 W.L. Jolly and K.D. Maguire, *Inorg. Synth.* 9 (1967) 102.
- 235 M.S. Haddad Thesis Americal University of Beerut (1975).
- 236 O. Glemser and H. Perl, *Naturwisson.* 48 (1961) 620.
- 237 S.A. Lipp and W.L. Jolly, *Inorg. Chem.* 10 (1971) 33.
- 238 H. Vincent and Y. Monteil, *Synth. React. Inorg. Metal. Org. Chem.* 8 (1958) 51.
- 239 R.L. Dekock and M.S. Haddad, *Inorg. Chem.* 16 (1977) 216.
- 240 N.R.D. Barton and A.W. Bubb, *J. Chem. Soc. Perkin. Trans.* (1977) 916.
- 241 W. Muthmann and E. Seitter, *Chem. Ber.* 30 (1877) 627.
- 242 A. Meuwsen, *Chem. Ber.* (1931) 2311.
- 243 P.A.G. O'Hare, J.L. Settle and W.M. Hubbard, *Trans. Faraday. Soc.* 62 (1966) 558.
- 244 R.L. Patton and W.L. Jolly, *Inorg. Chem.* 9 (1970) 1079.
- 245 M. Becke-Goehring and D. Schlafer, *Z. Naturforsch B21* (1966) 492.

- 246 A. Muller, N. Mohan, O. Glemser, S.J. Cyvin and N. Weinstock, J. Mol. Spectrosc. 59 (1976) 161.
- 247 H. Siebert, "Anwendender Schwingungs spekroskopie in der anorganischen chemie" Springer (1966).
- 248 A.D. Walsh, J. Chem. Soc. (1953) 2260.
- 249 N.K. Sanyal, R.K. Goel and A.N. Pandey, Spectrosc. Lett. 9 (1976) 681.
- 250 Ketelaar, Rec. Trav. Chem. 62 (1943) 289.
- 251 Ketelaar and Palmer, J. Am. Chem. Soc. 59 (1937) 9629.
- 252 Burns and Bernstein, J. Chem. Phys. 18 (1950) 1669.
- 253 Stephensen and Jones, J. Chem. Phys. 20 (1952) 135.
- 254 T. Beppu, E. Hirota and Y. Morino, J. Mol. Spectrosc. 36 (1970) 386.
- 255 R.L. Cook, J. Chem. Phys. 42 (1965) 2927.
- 256 T.L. Weatherly and O. Williams, J. Chem. Phys. 25 (1956) 717.
- 257 D.J. Millen and J. Pannel, J. Chem. Soc. (1961) 1322.
- 258 A.J. Banister, L.F. Moore and J.S. Padley, Spectrochim. Acta Part A23 (1967) 2705.
- 259 H. Garcil-Fernandez, C.R. Acad. Sci. C265 (1967) 88.
- 260 D. Neubauer and J. Weiss, Z. anorg. allgem. Chem. 340 (1965) 126.
- 261 M.G.B. Drew, D.H. Templeton and A. Zalkin. Inorg. Chem. 6 (1967) 1906.
- 262 R.C. Paul, C.L. Arora, J. Kishore and K.C. Malhotra, Aus. J. Chem. 24 (1971) 1637.
- 263 J. Weiss, Fortschr. Chem. Forsch, 5 (1966) 635.
- 264 P.J. Ashley and E.G. Torrible, Can. J. Chem. 47 (1969) 2587.
- 265 H. Wolbling, Z. Anorg. Allgem. Chem. 57 (1908) 286.
- 266 M. Becke Goehring and C. Magin, Z. Anorg. Allgem. Chem. 340 (1965) 126.

- 267 G.G. Alange and A.J. Banister, J. Inorg. Nucl. Chem. 40 (1978) 203.
- 268 M. Mecke-Goehring and G. Zirker, Z. Anorg. Allgem. Chem. 282 (1956) 83.
- 269 K.W. Daum, M. Becke Goehring and J. Weiss, Z. Anorg. Allgem. Chem. 278 (1955) 260.
- 270 M. Mecke Goehring and A. Debo, Z. Anorg. Allgem. Chem. 273 (1953) 319.
- 271 M. Beck Goehring, K.W. Daum and J. Weiss, Z. Naturforsch 101 (1955) 298.
- 272 E. Fluck, M. Goehring and J. Weiss, Z. Anorg. Allgem. Chem. 287 (1956) 51.
- 273 U. Thewalt, Angew. Chem. Int. Ed. Eng. 15 (1976) 765.
- 274 M. Goehring, J. Weiss and G. Zirker, Z. Anorg. Allgem. Chem. 278 (1955) 1.
- 275 O.C.M. Davis, J. Chem. Soc. 89 (1906) 1575.
- 276 J.A. Me Lean and A.G. Tumer, Inorg. Chim. Acta. 34(1) (1979) 199- 200.
- 277 M. Goehring and G. Zirker, Z. Anorg. allgem. Chem. 285 (1956) 70.
- 278 A. Meuwesen and M. Zosel, Z. Anorg. Allgem. Chem. 271 (1953) 221.
- 279 O. Glemser and H. Ludemann, Angew. Chem. 70 (1958) 190.
- 280 K.J. Wynne and W.L. Jolly, Inorg. Chem. 6 (1967) 107.
- 281 C.H. Chan and F.P. Olsen, Inorg. Chem. 11 (1972) 2836.
- 282 A.J. Banister and D. Younger, J. Inorg. Nucl. Chem. 32 (1970) 3763.
- 283 J.J. Zuckerman (Personal Communication).
- 284 K. Rotgers, Ph.D. Dissertation Berlin 1907.
- 285 B. Cohen, T.R. Hooper, D. Hugil and R.D. Peacock, Nature 207 (1965) 748.

- 286 O. Petersen and H.W. Roesky (Personal Communication).
- 287 M. Gochring, H. Hoherschutz and R. Appel, Z.Naturforsch B9 (1954) 678.
- 288 E.E. Aynsley and W.A. Campbell, J. Chem. Soc. (1957) 832.
- 289 A.J. Banister, J.A. Durrant I. Rayment and H.M.M. Shearer, J. Chem. Soc. Dalton Trans. (1976) 928.
- 290 R. Steudel, F. Rose, R. Reihardt, H. Bradaczek, Z.Naturforsch. 32B (1977) 488.
- 291 E.M. Holt, S.L. Holt, K.J. Watson, J. Chem. Soc. Dalton Trans. (1977) 514.
- 292 A. Bali, K.C. Malhotra, Ans. J. Chem. 29 (1976) 111.
- 293 J. Bojes and T. Chivers, J. Chem. Soc. Chem. Comm. (1977) 453.
- 294 J. Bojes and T. Chivers, J. Chem. Soc. Chem. Comm. (1978) 391.
- 295 J. Bojes and T. Chivers, Inorg. Chem. 17 (1978) 318.
- 296 J. Bojes, T. Chivers, I. Drummond and G. Maclean, Inorg. Chem. 17 (1978) 3668.
- 297 A. Meuwesen, Ber. 65 (1932) 1724.
- 298 M. Becke Goehring and D. Schuster, Z. Anorg. Allgem. Chem. 271 (1953) 281.
- 299 A. Meuwesen and O. Jakob, Z. Anorg. Allgem. Chem. 263 (1953) 200.
- 300 A.G. Mac Diarmid, J. Am. Chem. Soc. 78 (1956) 3871.
- 301 O. Glemser and E. Wyszomirski, Chem. Ber. 94 (1961) 1443.
- 302 R.F. Kruh, A.W. Cordes, R.M. Lawrence and R.G. Goforth, Acta. Crystal. 14 (1961) 1306.
- 303 M. Becke-Goehring and H.P. Latscha, Z. Naturforsch. 176 (1962) 125.
- 304 J. Weiss, Angew. Chem. 74 (1962) 216.
- 305 J. Weiss, Z. Anorg. Allgem. Chem. 333 (1964) 314.

- 306 A.W. Cordes, R.F. Kruh and E.K. Gordon, Inorg. Chem. 4 (1965) 681.
- 307 R.C. Paul, C.L. Arora and K.C. Malhotra, Chem. Ind. 51 (1968) 1810.
- 308 R.C. Paul, R.P. Sharma, S. Dogra, R.D. Verma, J. India. Chem. Soc. 53 (1976) 1228.
- 309 R.C. Paul, R.P. Sharma and R.D. Verma, Indian J. Chem. Soc.A. 15 (1977) 359.
- 310 R.D. Harcourt, J. Inorg. Nucl. Chem. 39 (1977) 237.
- 311 A.J. Banister, J.A. Durrant, J. Chem. Res.(9) (1978) 152.
- 312 R.J. Gillespie, D.R. Slim and J.D. Tyrer, J. Chem. Soc. Chem. Comm. (1977) 253.
- 313 J.G. Ballard, T. Birchall, R.J. Gillespie, E. Maharajh and D. Tyrer, Can. J. Chem. 56 (1978) 2417.
- 314 O.J. Scherer and G. Wolmerschäuser, Chem. Ber. 110 (1977) 3241.
- 315 T. Chivers and J. Proctor, Can. J. Chem. 57 (1979) 1286,  
Chem. Comm. 15 (1978) 642
- 316 A.J. Bamister, P.J. Dainh, A.C. Hazell, R.G. Hazell  
and J.G. Lambory, J. Chem. Soc. Chem. Comm. (1969) 1187.
- 317 R. Bartetzko and R. Gleiter, Inorg. Chem. 17 (1978) 995.
- 318 J. Chatt and J.R. Dilworth, J. Chem. Soc. Chem. Comm. (1974) 508.
- 319 M.W. Bishop, J. Chatt and J.R. Dilworth, J. Chem. Soc. Dalton Trans. 1 (1979) 1.
- 320 M.B. Hursthouse and M. Motevalli, J. Chem. Soc. Dalton Trans. (1979) 1362.
- 321 T.F. Brennen and I. Bernal, Inorg. Chim. Acta, 7 (1973) 28.
- 322 Special Publ. The Chem. Soc., London (1965) no. 18.
- 323 "Molecular Structures by Diffraction Methods" Vol. 2.  
Specialist Periodical Reports, The Chemical Society,
- 324 M.W. Bishop, J. Chatt, and J.R. Dilworth (unpublished work).



- 325 K.K. Pandey and U.C. Agarwala, Z. anorg. allgen. chem. 260(1) (1980) 000.
- 326 K.K. Pandey, R.C. Srivastava and U.C. Agarwala (unpublished result) (1980).
- 327 K.K. Pandey, S. Datta and U.C. Agarwala, Z. anorg. allgen. Chem. in press (1980).
- 328 K.C. Jain, K.K. Pandey, S.S. Katiyar and U.C. Agarwala, XXI ICCG, France (1980).
- 329 M.W. Bishop, S. Chatt and S.R. Dilworth, J. Chem. Soc. Chem. Comm. (1975) 780.
- 330 K.W. Muir, L. Manojlovic Muir and R. Herak (unpublished work).
- 331 K.K. Pandey and U.C. Agarwala, Angew. Chem. Int. Ed. Eng. in press (1980).
- 332 See Chapter II, Page 93, 94.
- 333 B.W.S. Kolthammer and P. Legzdins, J. Am. Chem. Soc. 100 (1978) 2247.
- 334 T.J. Greenhough, B.W.S. Kolthammer, P. Legzdins and J. Trotter, J. Chem. Soc. Chem. Comm. (1978) 1036.
- 335 T.J. Greenhough, B.W.S. Kolthammer, P. Legzdins and J. Trotter, Inorg. Chem. 18 (1979) 3548.
- 336 A.F. Berndt and R.E. Marsh, Acta. Cryst. 16 (1963) 118.
- 337 J.T. Malito (personal communication).
- 338 K.K. Pandey and U.C. Agarwala, Angew. Chem. Int. Ed. Eng. Communicated (1980).

### Infrared Spectra

Spectra of the compounds were recorded with a Parkin-  
Elmer 580 Infrared Diffraction Grating Spectrophotometer in  
the  $4000\text{--}200\text{ cm}^{-1}$  range. Samples were prepared as KBr and  
CsI pellets.

### Conductivity Measurements

Conductivity measurements of the compounds were made  
in the Elico conductivity meter type CM-80 of millimolar  
solutions in nitrobenzene.

### Magnetic Susceptibility Measurements

These measurements were made using a Gouy balance at  
room temperature ( $30^{\circ}\text{C}$ ).

### Melting Points

Melting points of the compounds were recorded on a  
Fisher-Johns melting point apparatus. The results are recorded  
in Table II.1.

### Results and Discussion

$(\text{NSCl}) (\text{solvent})_x$  reacts with  $\text{RhH}(\text{PPh}_3)_4$  to give a  
brown chlorobridged complex  $[\text{Rh}(\text{NS})\text{Cl}_2(\text{PPh}_3)]_2$  (I). The  
interaction of (I) with triphenylphosphine in  $\text{CH}_2\text{Cl}_2$  gave




Fig. 1 IR spectrum of  $[\text{Rh}(\text{NS})\text{Cl}_2(\text{PPh}_3)]_2$ .




Fig.2 IR spectrum of  $\text{Rh(NS)Cl}_2(\text{PPh}_3)_2$ .

a brown complex,  $\text{Rh}(\text{NS})\text{Cl}_2(\text{PPh}_3)_2$  (II). The infrared spectra of both the complexes I and II showed absorption bands  $1118\text{--}1120\text{ cm}^{-1}$  ( $\nu_{\text{NS}}$ ) and  $330\text{ cm}^{-1}$  (terminal  $\nu_{\text{Rh-Cl}}$ ) (Fig. 1 and 2). In the spectrum of I a band appeared at  $260\text{ cm}^{-1}$  which was not present in II. This band is assigned to bridged  $\nu_{\text{Rh-Cl}}$ . Both the complexes are non conducting in nitrobenzene, diamagnetic and air stable. The diamagnetism of complexes indicates the oxidation of rhodium is either +1 or +3. The values of  $\nu_{\text{NS}}$  ( $1120\text{ cm}^{-1}$ ) are in the range which has been reported for terminal bent coordinated thionitrosyl group [10].

Both the complexes I and II are soluble in benzene, dichloromethane, chloroform and insoluble in petroleum ether and hexane.

References

- 1 B.F.G. Johnson and J.A. Mc Cleverty, Prog. Inorg. Chem. 7, (1966) 277-359.
- 2 N.G. Connelly, Inorg. Chim. Acta Rev. 6, (1972) 47-89.
- 3 J. Chatt and J.R. Dilworth, J. Chem. Soc. Chem. Comm. (1974) 508.
- 4 M.W. Bishop, J. Chatt and J.R. Dilworth, J. Chem. Soc. Chem. Comm. (1975) 780.
- 5 M.W. Bishop, J. Chatt and J.R. Dilworth, J. Chem. Soc. Dalton Trans. 1 (1979), 1.
- 6 B.W.S. Kolthamer and P. Legzdins, J. Am. Chem. Soc. 100 (1978) 2247.
- 7 T.J. Greenhaugh, B.W.S. Kolthammer, P. Legzdins and J. Trotter, J. Chem. Soc. Chem. Comm. 23 (1978) 1036.
- 8 T.J. Greenhaugh, B.W.S. Kolthammer, P. Legzdins and J. Trotter, Inorg. Chem. 18 (1979) 3548.
- 9 K.K. Pandey and U.C. Agarwala, Z. Anorg. Allgen. Chem. 460 (1980) 000.
- 10 K.K. Pandey, S. Datta and U.C. Agarwala, Z. Anorg. Allgen. Chem. 465 (1980) 000.
- 11 K.K. Pandey and U.C. Agarwala, Angew. Chem. Int. Ed. Eng. in press (1980).
- 12 K.K. Pandey and U.C. Agarwala, Angew. Chem. Int. Ed. Eng. (Communicated) (1980).
- 13 K.C. Jain, K.K. Pandey, S.S. Katiyar and U.C. Agarwala, XXI ICCG (France) (1980).
- 14 N. Ahmad, J.J. Levison, S.D. Robinson and M.F. Uttley, Inorg. Synth. 15 (1974) 58.
- 15 W.L. Jolly, Inorg. Synth. 9 (1967) 102.

- 16 A.I. Vogel, 'Textbook of Quantitative Inorganic Analyses', 3rd Ed., Longmans Green, London (1961).
- 17 F.E. Beamish, 'A Textbook of Analytical Chemistry of Noble Metals', Oxford Pergamon Press (1966).

CHAPTER II

## THIONITROSYL COMPLEXES OF RHODIUM(III)

It has been known for many years that nitricoxide can coordinate to transition metals in a wide range of oxidation states to form metal nitrosyl $[1,2]$ . Analogous thionitrosyl complexes are markedly less abundant, a fact which reflects primarily the present lack of reagents which can be utilized to introduce the thionitrosyl group into a transition metal. Chatt, et.al. have reported the syntheses of thionitrosyl complexes of molybdenum, rhenium and osmium, containing linear thionitrosyl group $[3-5]$  and Legzdins, et.al. organometallic thionitrosyl complexes of chromium containing linear thionitrosyl group $[6-8]$  we have reported the thionitrosyl complexes of Ruthenium(II), Osmium(II), Rhodium(III) and Iridium(III), having three principal modes of bonding linear, bent or bridged, by the reaction of trithiazyltrichloride, which acts as a solvated monomer in THF or  $CCl_4-CHCl_3$   $[9-13]$ . This chapter describes the syntheses of thionitrosyl complexes of rhodium(III) by the reaction of hydridotetrakis (triphenylphosphine) rhodium(I) with trithiazyltrichloride in  $CCl_4-CHCl_3$  mixture. These compounds have been characterized by various physico chemical methods and their geometries have been proposed.




## EXPERIMENTAL

All the reagents used were of analar or chemically pure grade. All the solvents were dried and freshly distilled before use. The preparations were performed under the atmosphere of argon.

Hydridotetrakis(triphenylphosphine) rhodium(I) and trithiazyl-trichloride were prepared according to the methods described in the literature [14,15] .

### Preparation of Complexes



A yellow solution of  $(\text{NSCl})_3$  in  $\text{CCl}_4$  (20 ml) was added dropwise to a stirred solution of  $[\text{RhH}(\text{CO})(\text{PPh}_3)_3]$  (0.2g) in  $\text{CHCl}_3$  at room temperature. The mixture was stirred for 1h more followed by evaporation of solvent under reduced pressure. The residue was extracted in benzene. On adding petroleum ether to the benzene extract, a brown compound  $[\text{Rh}(\text{NS})\text{Cl}_2(\text{PPh}_3)]_2$  (m.p.  $215^\circ\text{C}$ ) was precipitated which was centrifuged out and washed with petroleum ether and water and dried in air. It was recrystallized from benzene-petroleum ether (1:3). Similar result was observed by using tetrahydrofuran in place of  $\text{CCl}_4$ - $\text{CHCl}_3$ .

Table II.1. Analytical data, melting points, colours and i.r. frequencies of the complexes

Compound	m.p. (°C)	Colour	Found		Analyses					Rh	P	I.R.frequencies (cm <sup>-1</sup> )	ν <sub>NS</sub>
			C	(calcd)	H	N	S	Cl					
[Rh(NS)Cl <sub>2</sub> (PPh <sub>3</sub> )] <sub>2</sub>	215	Brown	44.7 (44.8)		3.3 (3.1)	3.0 (2.9)	6.8 (6.6)	14.9 (14.7)	21.1 (21.3)	6.4 (6.4)		1118	
[Rh(NS)Cl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> ]	160- 163	Reddish brown	58.2 (58.0)		4.1 (4.0)	2.1 (1.9)	4.0 (4.3)	9.8 (9.5)	13.6 (13.8)	8.2 (8.3)		1120	